



Hydrogen sulfide (H₂S) - therapeutic relevance in rehabilitation and balneotherapy Systematic literature review and meta-analysis based on the PRISMA paradigm



MUNTEANU Constantin^{1,2,3}, MUNTEANU Diana^{1,5}, ONOSE Gelu^{2,4}

Editor: Mihail HOTETEU, Romanian Association of Balneology, hoteteu@yahoo.com



Balneo and PRM Research Journal DOI: <http://dx.doi.org/10.12680/balneo.2021.430> Vol.12, No.3 September 2021 p: 176-195

Corresponding author: *Constantin MUNTEANU*, E-mail: office@bioclima.ro

1. Romanian Association of Balneology, Bucharest, Romania
2. Teaching Emergency Hospital "Bagdasar-Arseni", Bucharest, Romania
3. Faculty of Medical Bioengineering, University of Medicine and Pharmacy "Grigore T. Popa", Iași, Romania
4. Faculty of Medicine, Department of Physical and Rehabilitation Medicine, University of Medicine and Pharmacy "Carol Davila", Bucharest,
5. National Institute of Rehabilitation, Physical Medicine and Balneoclimatology, Bucharest, Romania

Abstract

Background. An active molecule in sulfurous mineral - therapeutic waters and also in sapropelic mud is H₂S, a hormetic gaseous molecule that can actively penetrate the skin. While high levels of H₂S are extremely toxic, low levels are tolerated and have potential cytoprotective effects, with anti-inflammatory and antioxidant applications.

Objective. This systematic review aims to rigorously select related articles and identify within their content the main possible uses of hydrogen sulfide from balneary sources and to explain its physiological mechanisms and therapeutic properties.

Methods. To elaborate our systematic review, we have searched for relevant *open access* articles in 6 international databases: Cochrane¹, Elsevier¹, NCBI/PubMed¹, NCBI/PMC¹, PEDro¹, and ISI Web of Knowledge/Science¹, published from January 2016 until July 2021. The contextually quested keywords combinations/ syntaxes used are specified on this page. The eligible articles were analyzed in detail regarding pathologies addressed by hydrogen sulfide. All articles with any design (reviews, randomized controlled trials, non-randomized controlled trials, case-control studies, cross-sectional studies), if eligible according to the above-mentioned selection methodology, containing in the title the selected combinations, were included in the analysis. Articles were excluded in the second phase if they did not reach the relevance criterion.

Results. Our search identified, first, **291** articles. After eliminating the duplicates and non-ISI articles, remained **121** papers. In the second phase, we applied a PEDro selection filter, resulting in **108** articles that passed the relevance criterion and were included in this systematic review.

Conclusions. H₂S biology and medical relevance are not fully understood and used adequately for sanogenic or medical purposes. More research is needed to fully understand the mechanisms and importance of this therapeutic gas. The link between balneotherapy and medical rehabilitation regarding the usage of hydrogen sulfide emphasises the unity for this medical speciality.

Keywords: *Hydrogen sulfide / H₂S AND balneotherapy / inhalations / mud / Rheumatoid arthritis / neuroprotection / neurodegenerative disorders / Stroke / Parkinson / Alzheimer / Huntington / vascular dementia / Immune system / Diabetes / Cancer / Cardioprotection / Asthma / Alergic Rhinitis / COPD / COVID-19 / Burns / Analgesic effect / Clinical trials*

Authors:

Prof. dr. Gelu Onose, Ph.D.

Senior Physician in Physical and Rehabilitation Medicine (PRM) and Gerontology & Geriatrics (GG), Head of the P(neural-muscular)RM Discipline/ Clinic Division - the National Reference Centre for NeuroRehabilitation - and of its RDI Nucleus, and RDI (ex) Director, at the Teaching Emergency Hospital "Bagdasar-Arseni"



Assist. prof. dr. Constantin Munteanu, Ph.D.

Priority research areas: Bioclimatology, Hydrothermotherapy, and Physiology, President of the Romanian Association of Balneology, Vice President of the Romanian Society of Physical Medicine, Rehabilitation and Balneoclimatology,



Biol. Diana Munteanu, Priority research areas: microbiology, balneoclimatology, including the study of natural mineral waters and mofettes, therapeutic muds, and other balneary resources.



Reviewers:

Assist. prof. dr. Gabriela Dogaru, PhD,

Priority research areas: balneoclimatology, including the study of carbonated mineral waters and mofettes, molecular and cellular medicine, e-health systems, neurological and cardiovascular rehabilitation, development of spa, and rehabilitation tourism



Assoc. prof. dr. Mariana Rotariu, Ph.D. -

Psychomotricity, Kinesiology, psychomotor education and reeducation, Massage, Kinesitherapy, Kinesiprophilaxy, Complementary techniques, Occupational therapy, Rehabilitation, and techniques of functional reeducation



1. INTRODUCTION

Early life forms experienced and had to survive in an environment containing highly reactive chemicals such as NO, CO, and hydrogen sulfide (H₂S). During evolution, early life forms learned to tolerate these compounds and include them as essential molecules in their metabolism and signaling mechanisms.

Increasing evidence suggests that mammalian homeostasis strongly depends on a mutualistic relationship with gut bacteria. Interestingly, it has been found that metabolic and cardiovascular diseases, including hypertension, are associated with gut microbiota dysbiosis. Hydrogen sulfide (H₂S) regulates the functions of biological systems, including the circulatory system. H₂S released by bacteria in the colon may also contribute to the control of arterial blood pressure. Incidentally, sulfate-reducing bacteria are ubiquitous in the mammalian colon, and H₂S is just one among many molecules produced by the gut flora, and gut-bacteria-derived molecules, such as H₂S, have an important role in circulatory system homeostasis (1).

H₂S is, on the one hand, an endogenous gas signal molecule, which plays a vital role in various systems and diseases. H₂S is mainly produced in our body by three enzymes, including Cystathionine γ -lyase (CSE), Cystathionine β -synthase (CBS), and 3-mercaptopyruvate sulfurtransferase (3MST) together with cysteine aminotransferase (2). H₂S is also produced nonenzymatically from glucose, glutathione, thiosulfate, and sulfur-containing proteins and by the bacterial reduction of sulfur in the intestinal tract. H₂S is, on the other hand, an exogenous therapeutic gas, found in combination with carbon dioxide in sulfurous waters, with medicinal properties well known and appreciated in balneotherapy, and also in therapeutic muds (3).

Hydrogen sulfide (H₂S) naturally occurs in volcanic gases, natural gas, and some therapeutic waters (4). Also, H₂S is produced when bacteria break down organic matter in the absence of oxygen. H₂S is a flammable and colorless gas with a characteristic odor of rotten eggs. H₂S intoxication has been observed mainly in industrial environments. Workers may be exposed to H₂S in agriculture and industries, such as oil and wastewater treatment. H₂S is a toxic gas to humans, and acute exposure to large amounts of H₂S (> 500 ppm) can lead to death. The first biological experiment reported to study the effect of H₂S in animals was published in 1908 and described the lethal impacts of H₂S gas when absorbed through the skin or administered directly into the stomach or rectum. In the human oral microbiota, it has been found that H₂S is responsible for oral halitosis and is related to periodontal inflammation. In the intestinal microbiota (5), H₂S is a component of flatus (6).

The discovery of gaseous signaling molecules such as H₂S, NO, and CO has added a new era in biomedical

science, as these molecules are of great importance in mammalian physiology. They have been called "gas transmitters" because they are produced internally or synthesized (endogenously) in the body or are received from the atmosphere and transmit chemical signals promoting or inducing various physiological changes inside the mammalian body. Gas transmitters are permeable through the cell membranes, but their functions inside the body depend on their concentration.

Like other gas transmitters, the physiological role of H₂S has been overlooked or ignored due to its toxicity. The latest gas signaling molecule discovered after NO and CO has an enormous pathophysiological significance in various diseases and conditions. Due to its weak intermolecular strength, it is a sulfur analog of water and exists in gaseous form. It is synthesized both enzymatically and non-enzymatically within mammalian tissue, but the non-enzymatic pathway is of a more negligible significance.

Cystathionine β -synthase (CBS) and Cystathionine γ -lyase (CSE) are two enzymes responsible for H₂S biosynthesis from L-cysteine. Sulfurtransferase 3-mercaptopyruvate (3MST) is another enzyme that can generate H₂S via cys-catabolism. CSE and CBS are located in the cell's cytoplasm, but 3MST is partially expressed in the mitochondria and cytoplasm. A recent study showed that H₂S could be produced from D-cysteine by the enzyme D-amino acid oxidase (DAO). Non-enzymatically, H₂S can be produced from thiosulfate and glucose (by glycolysis) or phosphogluconate by NADPH oxidase (7).

Although H₂S has beneficial roles in various hematological diseases, urological diseases, cardiovascular function, and oxidative stress, the effects of H₂S in the CNS have attracted much more attention in recent years (8).

In vivo, H₂S can exert a dual effect on cell bioenergetics, at lower concentrations stimulating via sulfide - quinone oxidoreductase mitochondrial respiration and thus ATP synthesis or causing a reversible inhibition of cytochrome c oxidase at higher concentrations (9). H₂S has been shown to regulate all significant aspects of mitochondrial dynamics: mitochondrial fusion, mitochondrial fission, mitochondrial macroautophagy /mitophagy, and mitochondrial biogenesis). The majority of the published studies indicate that H₂S (especially in lower concentrations) tends to stabilize and preserve mitochondria, and, in many cases, can also stimulate mitochondrial biogenesis (10). The intricate interplay between H₂S and O₂ has been extensively investigated. As O₂ facilitates both the chemical and enzymatic oxidative decomposition of H₂S into persulfides and polysulfides, at low O₂ tension higher stability of H₂S is expected. Furthermore, hypoxic/ischemic conditions have been reported to enhance H₂S synthesis, through

upregulation or stimulation of the sulfide-synthesizing enzymes, accumulation of CBS in mitochondria, likely augmenting the H₂S mitochondrial levels, and release of CO-mediated inhibition of CBS and CSE. Hypoxia is thus expected to increase H₂S bioavailability, a condition that can have opposite physiological consequences. Indeed, while H₂S is protective against ischemic injuries, the enhanced biosynthesis and chemical stability of H₂S, combined with the reduced content in mitochondria, may increase the risk of H₂S toxicity in hypoxic cells (9).

Tissue development and repair have long been associated with sulfur metabolism (11). Since the first reports illustrating the expression of H₂S-producing enzymes in mammalian systems, there has been tremendous growth of preclinical studies investigating the protective effects of H₂S modulation, by way of donors or overexpression/inhibition of its synthesizing enzymes, against cardiovascular diseases. H₂S possesses infarct-sparing effects following myocardial I/R injury, but also it exerts anti-apoptotic, anti-oxidant, anti-inflammatory, and pro-angiogenic benefits under various cardiac stressors (12). The H₂S functions in the secretion of corticotrophin-releasing hormone from serotonergic neurons and the relaxation of smooth muscle. H₂S shields neurons and cardiac muscles from oxidative stresses and helps to maintain insulin secretion. H₂S has diverse physiologic functions such as relaxing blood vessels, lowering blood pressure, antiapoptosis, anti-inflammation, and anti-oxidative stress. Among these functions, the role of H₂S in antioxidative stress has been one of the main focuses over years (13).

2. METHOD

Literature Search Strategy. To elaborate our systematic review, we have searched for relevant *open access* articles, in 6 international databases: [Cochrane](https://www.cochrane.org/)¹, [Elsevier](https://www.elsevier.com/)², [NCBI/PubMed](https://pubmed.ncbi.nlm.nih.gov/)³, [NCBI/PMC](https://www.ncbi.nlm.nih.gov/pmc/)⁴, [PEDro](https://pedro.org.au/)⁵, and [ISI Web of Knowledge/Science](http://apps.webofknowledge.com/WOS_GeneralSearch)⁶, published from January 2016 until July 2021. The contextually quested keywords combinations/ syntaxes used were *Hydrogen sulfide / H₂S AND balneotherapy / inhalations / mud / Rheumatoid arthritis / neuroprotection / neurodegenerative disorders / Stroke / Parkinson / Alzheimer / Huntington / vascular dementia / Immune system / Diabetes / Cancer / Cardioprotection / Asthma / Alergic Rhinitis / COPD / COVID-19 / Burns / Analgesic effect / Clinical trials*. The eligible articles were analyzed in detail regarding pathologies addressed by hydrogen sulfide.

¹ <https://www.cochrane.org/>

² <https://www.elsevier.com/>

³ <https://pubmed.ncbi.nlm.nih.gov/>

⁴ <https://www.ncbi.nlm.nih.gov/pmc/>

⁵ <https://pedro.org.au/>

⁶ http://apps.webofknowledge.com/WOS_GeneralSearch

Inclusion and Exclusion Criteria

All articles with any design (reviews, randomized controlled trials, non-randomized controlled trials, case-control studies, cross-sectional studies), if eligible according to the above-mentioned selection methodology, containing in the title the selected combinations, were included in the analysis. Articles were excluded in the second phase if they did not reach the relevance criterion.

3. RESULTS

Our search identified, first, **291** articles. After eliminating the duplicates and non-ISI articles, remained **121** papers. In the second phase, we applied a PEDro selection filter, resulting in **108** articles that passed the relevance criterion and were included in this systematic review.




Fig. 1 PRISMA inspired flow diagram for articles search

4. A timeline of hydrogen sulfide research

The H₂S history – as an environmental toxin – dates back to 1700, with the observations of the Italian physician Bernardino Ramazzini (1633–1714), whose book “De Morbis Artificum Diatriba” is the reason why he is widely considered the founding father of the occupational medicine. Ramazzini predicted that this gas may damage the lungs (14). As a pure chemical species, the discovery of H₂S (1775) is credited to Swedish-German chemist Carl Wilhelm Scheele (1742– 1786), who produced it through the reaction of an acid with metal (poly)sulfides or by heating sulfur in hydrogen gas. In 1776, the chemical composition of the gas was determined by the French chemist Claude Louis Berthollet (1789).

The first biological experiment that directly investigated the effect of pure H₂S in animals was published in 1803 by French anatomist Francois Chaussier (1746–1828). He described the lethal effects of H₂S gas when the gas was absorbed through the skin. Subsequent studies used injections of H₂S-saturated solutions and reported respiratory effects (excitation, deep breathing), convulsive movements, and, at high doses, asphyxia and death. Over the subsequent 150 years, generations of toxicological researchers have investigated the

toxicological effects of H₂S in various species including humans. The results of these studies are now summarized in comprehensive monographs, including limits and methods of its detection, occupational exposure limits, and the multiple biological effects of H₂S on various cells and organs (14).

H₂S, at low concentrations, has an entirely different pharmacological profile than what was previously characterized in the toxicological literature (where, typically, high concentrations were used to demonstrate adverse/ noxious effects).

With the use of pharmacological inhibitors, as well as mice lacking various H₂S-producing enzymes, and by using transient and stable silencing of H₂S-producing enzymes, the functional role of endogenously produced H₂S began to emerge in more detail (as opposed to testing the effect of low concentrations of exogenously applied H₂S on biological systems). There were also significant advances made in the field of H₂S-mediated signaling, including the concept of sulfhydration (post-translational modification of protein cysteine residues, with functional consequences), which is prominently induced by polysulfides. Another area within the growing field of H₂S biology relates to the interactions of H₂S with various oxygen- and nitrogen-derived species, which includes functional interactions as well as the recognition of the biological roles of various hybrid S/N species. The state-of-the-art of H₂S biology can be overviewed in general reviews and monographs as well as in specialized review articles focusing on the roles of H₂S in the regulation of the nervous system, cardiovascular system, gastrointestinal system, renal system, metabolic and mitochondrial aspects, cellular signaling, and the biochemistry of the various H₂S-producing enzymes. Separate articles focus on the details of the chemical properties and reactivity of H₂S and its functional interactions with other reactive species including oxygen-derived oxidants and free radicals and NO. Other review articles focus on pharmacological donors and inhibitors of H₂S biosynthesis, and the therapeutic aspects and translational potential of H₂S biology (14).

5. Chemical and Physical Properties of H₂S

Aqueous species elements can be quantified by chemical speciation analysis or estimated by speciation modeling based on chemical thermodynamics (15). Although H₂S exists as a gaseous molecule under ambient temperature and pressure, it can easily dissolve in water with a solubility of around 80 mM at 37 °C. The half-life of H₂S in air varies from 12 to 37 h (16). Upon dissolution, H₂S promptly reaches the equilibrium of H₂S /HS⁻/S²⁻ species, which is different from the other two gaseous transmitters, respectively, NO and CO. Based on its PKa value, it is estimated that there will be 14% of H₂S, 86% of HS⁻, and a negligible level of S²⁻ in physiological solution (pH 7.4, 140 mM NaCl, 37 °C).

Moreover, H₂S gas is highly lipophilic, which allows its free penetration into biological membranes of all types and becomes biologically active. Sulfur is a versatile atom with multiple oxidation states ranging from -2 to +6. In H₂S, the oxidation state of sulfur is -2; therefore, it can act as a reductant and only be oxidized. In line with this, accumulative evidence has suggested that H₂S actively takes part in three reactions in mammalian systems, including interaction with the reactive oxygen species (ROS) or reactive nitrogen species (RNS) species; reduction of the protein irons, namely, the metal center on proteins; transformation of protein R-SH group into R-SSH group, a process called S-sulfhydration (17).

6. H₂S toxic effects

There are more than 300 years since the first description of H₂S as a poisonous molecule. Among the more recent studies, the work of Attene-Ramos should be mentioned, who demonstrated the genotoxic effect of high doses of H₂S, Nicholson, Khan, and later Dorman and colleagues have directly demonstrated the inhibition of cytochrome c oxidase activity *ex vivo* in tissues after H₂S exposure of experimental animals and implicating these effects in the disruption of respiratory and mitochondrial functions in the mammalian brain (and other tissues) after H₂S exposure *in vivo*. H₂S has wide-ranging effects on gene expression (associated with a variety of biological processes including cell cycle regulation, cellular division, DNA metabolism and repair, protein kinase regulation, cytoskeletal organization, and biogenesis). It is currently accepted that H₂S exerts its toxicological actions on the molecular level primarily through the inhibition of mitochondrial respiration via inhibition of mitochondrial Complex IV (cytochrome c oxidase). Via this action, the consumption of O₂ is inhibited, and electron transport and ATP generation are blocked (14).

7. The physiological role of hydrogen sulfide

Under physiological conditions (aqueous solutions at pH 7.4), one-third of H₂S is un-dissociated and two-thirds is hydrolyzed to sulfide and hydrosulfide ions in the following reaction: $H_2S \leftrightarrow H^+ + HS^- \leftrightarrow 2H^+ + S^{2-}$. In mammalian tissues and blood, the concentration of H₂S is 1–160 μM, under physiological conditions. Higher concentrations of H₂S are present in the brain (50–160 μM) and blood (10–100 μM) (18).

Through the participation of H₂S in many physiological and pathological processes, including its role in regulating inflammatory processes, hypoxia, cell proliferation, apoptosis, neuromodulation, and cardioprotection, H₂S is now accepted as a player - as a gas transmitter (gas signaling compound) that is equally important as nitric oxide and carbon monoxide in mammals and as a signaling molecule as necessary as hydrogen peroxide (H₂O₂) in plants.

Hydrogen sulfide (H₂S) has emerged as a critical mediator of multiple physiological processes in

mammalian systems. The pathways involved in the production, consumption, and mechanism of action of H₂S appear to be sensitive to alterations in the cellular redox state and O₂ tension. Indeed, the catabolism of H₂S through a putative oxidation pathway, the sulfide quinone oxidoreductase system, is highly dependent on O₂ tension. Dysregulation of H₂S homeostasis has also been implicated in numerous pathological conditions and diseases (19).

8. Measurement methods of H₂S

Reliable bioanalytical methods for highly specific and sensitive detection of H₂S are key for the investigation of H₂S functions in biological systems. Over the past few decades, several techniques have been reported for H₂S detection in bulk solutions. Of various approaches, optical detection using H₂S responsive sensors has been recognized as one of the most promising technology due to its inherent advantages, such as high sensitivity and simplicity, rapidity, and efficiency. The H₂S responsive sensors are normally designed based on different sensing mechanisms, such as thiolysis of dinitrophenyl ether, reaction with azide, nucleophilic addition with formaldehyde groups, and displacement of metal ions from luminescent dyes. Under these reactions, several sensors have been developed for sensitive and selective H₂S detection and H₂S visualization in live cells and organisms. The majority of the reported H₂S sensors are designed based on the “OFF-ON” changes of luminescent signals (20).

H₂S in aqueous solution can be found in the form of hydrogen sulfide gas (H₂S) or one of its dissociated forms, hydrosulfide anions (HS⁻) and sulfide anions (S²⁻), although at physiological pH, S²⁻ is found only in a negligible concentration. In addition, H₂S can bind to some biological matrices (proteins, glutathione, etc.) and can dissociate in response to a physiological stimulus in free H₂S. Moreover, the HS and H₂S anion have a high tendency to oxidize, especially in the presence of metal ions and oxygen in aqueous solutions (21).

The first H₂S quantifier was published in 1949 and included spectrophotometric determination using the methylene blue method, but the sensitivity was very low. In 1969, an improved method based on the methylene blue method using *n,n*-dimethyl- β -phenylenediamine sulfate was described, which increased the sensitivity of the method by 10%, with concentration limits of 1 to 1000 μ M. However, the main disadvantages of this method for biological sample measurements are low sensitivity, overestimation of H₂S below acidic pH, and interference due to turbidity of biological samples. Monobromobiman (MBB) derivative: this is another method widely used in biological samples to measure H₂S concentration. In this method, H₂S is derivatized into a sulfide-dibiman product, which HPLC can subsequently measure due to its fluorescence. MBB sensitivity reaches

the nanomolar range, but the weaknesses of this method are the instability of the standards and the need for a complete pH control when comparing the samples. Another advantage of this method is that it allows the detection and quantification of all three biological forms of sulfur: free hydrogen sulfide, labile acid sulfur, and bound sulfan sulfur. An improved MBB-based method was developed that included ³⁵S-labeled sulfide-dibiman as an internal standard and measured the derivatized products by liquid chromatography-mass spectrometry.

9. H₂S Mechanisms of Action

There is an important effect of H₂S binding to target proteins such as cytochrome c oxidase, hemoglobin, and myoglobin, among others. It has, however, become widely accepted that a huge number of the processes controlled by H₂S are caused by a posttranslational modification of cysteine residues called persulfidation. Protein persulfidation is an oxidative posttranslational modification of cysteine residues caused by H₂S, in which a thiol group (-SH) is transformed to a persulfide group (-SSH). Sulfane sulfur species, persulfides, and polysulfides are more nucleophilic than H₂S (22).

It is well established that H₂S regulates different physiological processes in cells directly or by crosstalk with other signaling molecules. There is clear evidence of crosstalk of H₂S and NO in the literature. In mammals, both gasotransmitters interact with each other to modulate the cardiovascular system by regulating angiogenesis and endothelium-dependent vasorelaxation, and to modulate Alzheimer's disease by regulating pathways involved in the central nervous system. Furthermore, inhibition of NO generation by H₂S has been extensively studied, but there is also evidence that NO can activate the production of H₂S in endothelial cells. However, NO can bind to cystathionine β -synthase (CBS), which is responsible for H₂S biosynthesis and can impede its enzymatic activity, showing the complexity of the crosstalk between these two gasotransmitters.

Carbon monoxide is another important gasotransmitter in animals; carbon monoxide is generated from oxidative degradation of heme by the enzyme heme oxygenase. CO may inhibit CBS activity and therefore may modulate H₂S biosynthesis. Exogenous H₂S also upregulates the CO/heme oxygenase system in the pulmonary arteries of hypoxic rats and stimulates heme oxygenase levels.

H₂S is a regulator of glucose homeostasis and plays an important role in the metabolism of hormones, such as insulin and glucagon. It has been demonstrated that β cells of the pancreas can produce high levels of H₂S, predominantly by cystathionine γ -lyase (CSE), which blocks glucose-stimulated insulin secretion. This effect is caused by increased endoplasmic reticulum stress, leading to apoptosis of β cells, which drives the reduction in insulin secretion. Some other studies revealed the importance of H₂S in the modulation of estrogen receptor

expression and its anti-proliferative effect on vascular smooth muscle cell growth and proliferation. Further research concluded that the antiatherosclerotic effect of estrogen is mediated by CSE-generated H₂S and that H₂S production in the liver and vascular tissues is enhanced by estrogen via its stimulatory effect on CSE activity. H₂S signaling was also linked with the regulation of two endocrine hormones associated with longevity control, growth hormone and thyroid hormone.

10. Hydrogen sulfide (H₂S) role in Balneotherapy

Balneotherapy (Latin *balneum*, bath (23)) has been defined as the use of natural mineral waters (24), natural peloids and mud, and natural sources of different gases (CO₂, H₂S, and Rn) for medical purposes such as prevention, treatment, and rehabilitation (25)

Among the inorganic molecules, which generally constitute the mineral waters, sulfur has currently been recognized as a crucial element with a wide range of functions, mainly when it was found in the form of hydrogen sulfide (H₂S). Sulfurous waters (26) contain at least 1 mg of H₂S, HS, S, or thiosulphates per liter or sulfur colloidal complexes, in the form of simple or mixed sulfurous waters (alkaline earth, carbonated, sodium chlorinated). H₂S represents the main active molecule of sulfurous mineral-medicinal waters (27). It has been also shown by *in vitro* and *in vivo* studies that there is a positive action of mineral waters on the oxidant/antioxidant system. Many authors investigated the possible activity of different hydrogen sulfide (H₂S) donors in normal or osteoarthritic (28) chondrocytes stimulated by proinflammatory stimuli such as Interleukin-1b or lipopolysaccharide. The incubation with exogenous H₂S sources inhibits the release of Nitric oxide, Prostaglandin E₂, Metalloproteases, Tumor necrosis Factor- α , Interleukin-6, Interleukin-8 and decreases the Nuclear Factor-kB activation (29). It is well known that sulfurous water baths were used by ancient civilizations and were known to have healing effects against particular diseases. H₂S has been recognized as having anti-inflammatory, anti-bacterial, vasodilator, and anti-fungal properties owing to its sulfur content. Sulphur (S), Hydrogen Sulphide (H₂S), Sulphate (SO₄²⁻) are involved in cartilage, hair/nails formation, enzyme activity in redox processes, and cellular respiration. SO₄²⁻ is essential to several metabolic and cellular processes, particularly in fetal growth and development (30).

Sapropel's complex chemical and biological structure explains its multifunctional effect on the body (31). Baths stimulate redox reactions, which is indicated by an increase in the hemoglobin level and erythrocyte count. observed also in patients with rheumatoid arthritis (32) receiving artificial sulfur and hydrogen sulfide baths (33). The potential of sulfurous thermal water (STWs) to directly regulate bone cells function and to use STWs in

complementary medicine for the treatment of bone-wasting diseases is a new terrain of investigation. A growing body of evidence shows that H₂S has therapeutic potential in age-associated diseases. Exogenous H₂S stimulation induces osteogenic differentiation. H₂S donors can be used, similarly to other osteoinductive molecules, to increase the osteogenicity of bone implants, and most importantly that therapeutic intraperitoneal administration of H₂S prevents the onset of bone loss in the mouse model of osteoporosis (34). Hydrogen sulfide (H₂S) seems to affect cells involved in degenerative joint diseases, it was the objective of this study to investigate the effects of exogenous H₂S on fibroblast-like synoviocytes, which are key players in OA pathogenesis being capable of producing pro-inflammatory cytokines and matrix-degrading enzymes (35).

Healthy cartilage maintenance relies on an equilibrium among the anabolic and catabolic processes in chondrocytes. Hydrogen sulfide (H₂S) has shown anti-inflammatory and anti-catabolic properties in several *in vitro* and *in vivo* models. H₂S in the musculoskeletal system is quite incipient. Yet, it is already accepted that it is present and modulated in the osteoarthritic process and it exerts anti-inflammatory and cytoprotective effects in joint cells at low micromolar doses. H₂S is present in the synovial fluid (SF). In patients with rheumatoid arthritis, SF H₂S levels were significantly and substantially higher than in paired plasma samples or SF aspirates from OA patients. Additionally, H₂S levels in SF and plasma correlated with clinical markers of disease activity and inflammation (36).

Bone tissue is constantly remodeling. It responds to various stimuli, such as physical exercise. H₂S can prevent the onset of bone loss in the mouse model of osteoporosis (37).

11. Balneo-rehabilitative role of H₂S in Romania

In our country, sulfurous waters have special healing qualities. One of the most important sources of sulfurous water is found in Dâmbovița County, in Pucioasa. On the Olt valley, at Calimanesti, Caciulata, the sulfurous waters, through the different concentrations in hydrogen sulfide, are used therapeutically both in the internal and external cure. The localities of Baile Govora and Harghita Bai are also famous, but also Baile Bârzava, Baile Boghiș, Olanesti and those on the sea coast, Venus and Mangalia. In Herculane, a resort arranged and operated for the first time 2000 years ago, it benefits from the sulfurous waters of special quality and the possibility of performing thermal bath cures. Sulfurous waters are indicated both for crenotherapy and in external treatment in the form of baths, inhalations, and vaginal irrigation, due to the effects of H₂S, which is reabsorbed through the skin and gastric mucosa, upper airways, bronchopulmonary and vaginal mucosa.

Diseases that can be treated with sulfurous water:

1. Crenotherapy with sulfurous waters, usually mixed, has effects of stimulating gastric secretion as well as intestinal peristalsis, choleric and cholagogue effects on the liver and bile ducts, antitoxic effects, and lowering blood sugar in diabetics. The mixed carbonated ones also have diuretic effects through the action of CO₂. The indications for crenotherapy with low concentration sulfurous waters are hepato-biliary disorders, especially biliary dyskinesias, gallstones, and chronic cholecystitis, urinary tract infections, forms of diabetes, and heavy metal poisoning.
2. Diseases of the musculoskeletal system. These diseases include rheumatic, inflammatory, degenerative, abarticular diseases, post-traumatic sequelae of the hands and feet, etc.
3. Peripheral neurological disorders. Post-traumatic paralysis and parasites of the limbs; polyneuropathy after the acute phase; ponytail syndrome; sequelae after polio with secondary tendon damage, for corrective surgery.
4. Gynecological diseases. Sulfur waters also have an anti-inflammatory (38) and anti-allergic effect in gynecological diseases, favorably influencing the secretory glands of the cervix. It also promotes the restoration of the intravaginal biological component.
5. Respiratory disorders. The treatment for prophylactic purposes is addressed to workers who work with dust, lint, toxic gases, humidity or dry air, temperature variations, etc., and who are prone to inflammatory or allergic bronchopulmonary diseases. Treatment is recommended for convalescents after acute, bacterial and viral pneumonia, simple chronic tracheobronchitis, respiratory neurosis. The sulfurous mineral waters used in inhalation form have an action of wetting the bronchial trajectory, favoring the drainage of the products of bronchial secretion.

12. H₂S role on neurodegenerative disorders

In some brain disorders, endogenous H₂S generation is blunted, and H₂S deficiency has been implicated in the progression of Alzheimer's disease (AD), Parkinson's disease (PD), Huntington's disease (HD), and traumatic brain injury (TBI). Although the mechanisms of action of H₂S have not been fully characterized, redox regulation has been identified as a critical player in H₂S-mediated neuroprotection and toxicity (39).

H₂S has a significant neuroprotective role in the central nervous system. H₂S provided neuroprotection in TBI via inhibition of autophagy and apoptosis (40). H₂S exhibits cytoprotection against injuries in many tissues/organs through its anti-oxidative, anti-inflammatory, and/or anti-apoptotic effects. Neuroprotection of exogenous H₂S against cerebral injury is associated with the reduced accumulation of autophagic vacuoles (41).

Important signaling events of H₂S in various neuronal cells/cell lines are listed below:

1. Inhibition of monoamine oxidase (via catecholamines)
 2. NMDA potentiation (via glutamate)
 3. Cystic fibrosis transmembrane conductance regulator (CTFR) channel activation (via chloride channels)
 4. KATP and KCa²⁺ channel activation (via potassium channels)
 5. Intracellular calcium mobilization, L-type and T-type channel activation (via calcium channels)
 6. Suppression of various types of neuronal toxicity (via oxidative stress)
 7. Inhibition of p38-MAPK (via mitogen and tyrosine kinase receptors)
 8. Stimulation of PKA and elevation of cAMP (via PKA)
- AD, a common form of dementia, characterized by memory impairment, personality changes, and various neuropsychiatric symptoms which cause neuronal apoptosis, neuronal inflammation (induced by amyloid-β), and increased oxidative stress.

The level of H₂S in the brain of a patient with AD is lower than healthy people of the same age. A recent study revealed that in a rat model of vascular dementia, plasma H₂S level was lower and i.p. injection of NaHS (H₂S donor) protected neuronal injury and improved behavioral (learning and memory) test results. Another study demonstrated that progression of AD was abruptly after treatment with spa-water rich in H₂S content. The role of H₂S in the improvement of cognitive functioning, spatial learning and memory, and neuroprotective effects is also providing us hopes against AD.

Research and studies regarding H₂S are still in the preliminary stage in the field of biomedical science. Further research also should be focused on their combinatorial effect and signaling pathways regarding their antagonistic effect should be disclosed (8).

13. Hydrogen sulfide and Stroke

Stroke (ischemic or hemorrhagic) is an acute cerebrovascular disease, a leading cause of death and disability worldwide (42). H₂S has tissues concentrations of 50-160 mM, much higher than in peripheral blood (0-46 mM), suggesting that H₂S may have some effects and potential therapeutic value (43). H₂S can facilitate hippocampal long-term potentiation *via* activating the N-methyl-D-aspartic acid (NMDA) receptor. H₂S can activate calcium channels to increase calcium influx, whereas decreasing the release of calcium stored in the cells, thus induce the production of calcium waves in primary cultures of astrocytes. This process can be inhibited by a calcium channel antagonist. In addition, H₂S has been known as ATP-sensitive potassium (KATP) channel activator in secondary brain injury after stroke or other diseases, H₂S could play its neuroprotection through mimicking the KATP channel (42).

Fibrinolytic therapy with recombinant tissue plasminogen activator (tPA) is a globally accepted treatment for ischemic stroke. Unfortunately, the clinical application of tPA remains rather limited due to the enhanced risk of cerebral hemorrhage associated with tPA therapy. H₂S-base therapy represents a novel approach to increase the safety of tPA thrombolysis in stroke patients. H₂S donor inhibited the Akt-VEGF-MMP9 cascade in both the mouse MCAO model and the endothelial OGD model following tPA treatment (44). Given the ability of GSH to function as a histone modifier, it would be appropriate to consider H₂S as an epigenetic modulator albeit in an indirect manner. Concerning oxidative stress, H₂S is also mitoprotective as it averts energy failure and upregulates superoxide dismutase (SOD) expression with a reciprocal downregulation of cytochrome oxidase. Paradoxically, both inhibition of H₂S production and therapeutic H₂S donation seem to substantially influence the disease progression in various animal models of inflammation, reperfusion injury, and circulatory shock.

Treatment with H₂S also has been shown to reduce brain injuries and post-ischemic cerebral edema in a dose-dependent manner possibly by blocking apoptosis. (45).

H₂S also regulates the cerebral microvascular response to hypoxia. Some reports showed that under normal O₂ conditions, CO binds to the heme in CBS and inhibits the activity of producing H₂S. Hypoxia induces the production of H₂S through the decline of the generation of CO and vasodilation. H₂S has been reported to modify specific cysteine residues in proteins through the formation of a polysulfide and persulfide bond and to be involved in the regulation of such activity by sulfhydrylation of the enzyme (46). H₂S has an important role in maintaining the stability of the cerebrovascular environment. The decrease of H₂S level leads to endothelial dysfunction and promotes the development of ischemic stroke (47).

14. Hydrogen sulfide and Parkinson disease

Parkinson's disease (PD) is the second most common neurodegenerative disorder affecting over four million people worldwide, of high prevalence among people elder than 65 years of age (48). The brain of a PD patient is featured with the loss of dopamine (DA) - secreting neurons in an area of the midbrain called the substantia nigra, therefore causing bradykinesia, postural instability, resting tremor, and rigidity of patients (49). Organ-protective effects of inhaled H₂S have generally been attributed to its ability to induce hypothermia in rodents. Deep hypothermia induced by H₂S breathing reduced cerebral-infarct volume in rats after focal ischemia (49). Oxidative damage is considered a primary pathogenic mechanism of nigral dopaminergic cell death in PD (50). Despite decades of efforts, medications to cure or slow the progress of PD are still lacking. Emerging evidence suggests that H₂S maybe not only be implicated in the

pathogenesis of PD but also protect neuronal loss in various PD models, suggesting that this gaseous molecule may be new hope for the devastating disease (17).

The presently clinically used drug for treating Parkinson's disease is L-DOPA (51), which only ameliorates the symptoms but could not reverse the process of DA neuronal degeneration (52). A reduction in endogenous H₂S in the striatum appears to be responsible for the onset of Parkinson's disease and that exogenous H₂S treatment results in attenuation of DA neuronal degeneration in a mouse model of Parkinson's disease. These findings suggest a neuroprotective effect of H₂S during the neural injury of Parkinson's disease. Hydrogen sulfide appears to confer cytoprotection via multiple mechanisms including antioxidant and antiapoptotic effects. H₂S may upregulate endogenous antioxidants through a nuclear-factor-E2-related factor-2 (Nrf2)-dependent signaling pathway. Nrf2 regulates gene expression of several antioxidant proteins and phase II detoxification enzymes (for example, glutathione S-transferase (GST)). Whether H₂S protects neurons by upregulating Nrf2-dependent antioxidant mechanisms remains to be determined. The role of the Rho/ROCK/LIMK pathway in neurodegeneration has been acknowledged in recent studies. Activated ROCK2 is significantly inhibited by H₂S. It has been well established that microglial ROCK leads to phagocytosis of dopaminergic neurons. In addition, CaMKK-beta-dependent AMPK activation causes NaHS suppression of neuroinflammation (48).

Adult neurogenesis is one method to combat the loss of DA neurons in the adult midbrain, which may prevent cell death and relieve dyskinesia in PD. Previous studies have focused on DA neurons and how to delay neuronal degeneration by treatment with H₂S. However, it remains unclear whether H₂S therapy could induce neurogenesis in adults to alleviate the progression of PD (53).

15. Hydrogen sulfide and Alzheimer disease

Alzheimer's disease (AD), the most common cause of dementia and neurodegeneration in the elderly, is characterized by deterioration of memory and executive and motor functions (54). Neuropathologic hallmarks of AD include neurofibrillary tangles (NFTs), paired helical filaments, and amyloid plaques. Mutations in the microtubule-associated protein Tau, a major component of the NFTs, cause its hyperphosphorylation in AD. Signaling by the gaseous molecule hydrogen sulfide (H₂S) is dysregulated during aging. H₂S signals via a posttranslational modification termed sulfhydrylation / persulfidation, which participates in diverse cellular processes. Cystathionine γ -lyase (CSE), binds wild-type Tau, which enhances its catalytic activity. By contrast, CSE fails to bind Tau P301L, a mutant that is present in the 3xTg-AD mouse model of AD (55).

Hydrogen sulfide plays an important role in vasorelaxation, inflammation, cell angiogenesis, hippocampal memory formation, and cellular bioenergetics. In addition, it was observed that H₂S levels were decreased, and Hcy levels were elevated in AD brains. Likewise, it was noted that the physiological concentration of H₂S specifically enhances the NMDA receptor-mediated response and facilitates hippocampal long-term potentiation (LTP), which is extensively associated with proper learning and memory function.

Matrix metalloproteinases 9 (MMP9) role is well known in the regulation of synapse function and plasticity, its association with Hcy-induced functional remodeling of synapses remains uncertain. Administration of Hcy into the mice brain affects memory function and H₂S seems to protect against Hcy-induced neurotoxicity, but the underlying pathology was unclear. The therapeutic potential of H₂S on synaptic remodeling. Additionally, these data may be critical to better understand the role of MMPs in neurovascular injury in hyperhomocysteinemia. However, more detailed studies using this model may be warranted to more closely simulate the clinical scenario in cerebrovascular diseases such as Alzheimer's and stroke (56).

The level of H₂S is decreased in AD patients and that this change in the H₂S levels may be related to the severity of AD (57). The emerging mitochondrial roles of H₂S include antioxidant, antiapoptotic, and anti-inflammatory effects. H₂S modulates A β -induced damage attenuating the increase in intracellular ROS levels. H₂S may attenuate spatial memory impairment and neuroinflammation within the hippocampus of AD model mice. With regard to the regulatory role of H₂S in cellular bioenergy metabolism, recent studies have shown that H₂S can serve as a physiological electron donor and as an inorganic energy source in mammalian cells. H₂S on mitochondrial dysfunction in AD models are significant (54).

16. Hydrogen sulfide and Huntington disease

Huntington's disease (HD) is an autosomal dominant neurodegenerative disease caused by the expansion of CAG repeats in the gene huntingtin, htt, present on chromosome 4. About 1 in 7500 individuals are affected by HD worldwide, and no satisfactory cure exists so far. HD primarily affects the corpus striatum of the brain and manifests as abnormal involuntary movements, motor and cognitive deficits. Once symptoms appear, median survival is about 18 years. The number of CAG repeats in mutant htt (mHtt) inversely correlates with the age of onset of disease. Translation of the CAG repeats results in an abnormally long polyglutamine repeat at the N-terminal end of Htt. mHtt is proteolytically cleaved and the N-terminal fragments aggregate to form inclusion bodies, a characteristic feature of the disease, although its role in disease progression is debated. Immunostaining

with antibodies against mHtt led to the discovery of intranuclear inclusions in mouse models and patient samples, a hallmark of the disease. mHtt impacts multiple cellular processes ranging from transcriptional and translational regulation, mitochondrial function, DNA replication and repair to nucleocytoplasmic transport which leads to neurotoxicity (58).

H₂S signals *via* sulfhydration, wherein it mediates the conversion of -SH groups of reactive cysteine residues on target proteins to -SSH or persulfide groups. H₂S is synthesized by cystathionine gamma lyase (CSE), cystathionine beta synthase (CBS) and 3-mercaptopyruvate sulfurtransferase of the reverse transsulfuration pathway. Aberrant H₂S metabolism is involved in the progressive neurodegeneration seen in Huntington's disease (HD). The neuronal redox effector functions of H₂S are differentially modulated by the action of mHtt on CSE and CBS thereby affecting disease progression in HD by variable regulatory mechanisms.

17. Hydrogen sulfide and vascular dementia

Epidemiological studies show that vascular dementia (VD) is the second commonest cause of dementia after Alzheimer's disease (AD), and results from ischemic, ischemic-hypoxic, or hemorrhagic brain lesions.

In CNS, studies suggest that Notch signaling regulates neuronal synaptic plasticity, and therefore learning and memory. Interestingly, Notch signaling pathway has crosstalk with other signaling pathways, such as Akt, Wnt, and NF- κ B pathways. Studies report that GSK-3 β modulates Notch signaling. The production of H₂S is high in the brain, considered as a neuromodulator. A previous study demonstrate that H₂S facilitates the induction of hippocampal LTP by enhancing the activity of NMDARs in neurons. H₂S improves LTP in hippocampus and inhibits the neuronal death induced by bilateral common carotid arteries ligation. H₂S attenuates neuronal injury induced by VD via inhibiting apoptosis (59).

18. Hydrogen sulfide and the Immune system

Concerning cell-mediated immunity, balneotherapy applications can improve the survival capacity and activity of neutrophils. Sulphurous mineral water favours the short-term survival of neutrophils, speeding up the resolution of infections and preventing further inflammation. Studies conducted in patients with osteoarthritis who underwent hydrotherapy or mud-bath therapy showed that, after the treatment, neutrophils' circulating and functional capacity increased, reflecting a greater defence capacity against pathogens and thus a potential lower susceptibility to infections. Balneotherapy may also contribute to increase cortisol levels in healthy and subhealthy people. Changes in cortisol levels suggest that mineral baths may modulate the activity of the hypothalamic-pituitary-adrenal axis, inducing a transiently but significant rise in ACTH production (60).

One important property of H₂S is its bell-shaped (or biphasic/bimodal) pharmacological mode of action: in many conditions, low concentrations of H₂S can exert physiological, regulatory or modulatory effects, and act as cytoprotective, antioxidant and anti-inflammatory agents. Neutrophils are the most abundant leukocytes in the blood and constitute the first line of immune defense against numerous infectious pathogens (bacteria, fungi, protozoa). They are traditionally viewed primarily as phagocytic cells and are among the first to migrate to infected sites for pathogen killing and debris removal. Neutrophil-mediated killing of invading pathogens involves several different strategies: engulfment of microbes, secretion of anti-microbials and formation of neutrophil extracellular traps (NETs). The inhibitory effect of endogenous and exogenous H₂S on neutrophil adhesion has been confirmed in several studies. monocytes/macrophages express all 3 major H₂S-producing enzymes, with perhaps CSE and 3-MST playing the major roles. The role of H₂S in the regulation of macrophage chemotaxis is complex. The fact that H₂S affects multiple signalling pathways and immunological and inflammatory mechanisms can be a strength from a translational standpoint: simultaneously downregulating multiple interacting pathways of inflammation (61) or cell death may more effective than targeting a single pathway that could be bypassed or replaced by various compensatory mechanisms (62).

19. Hydrogen sulfide and Diabetes

Diabetes mellitus is one of the most prevalent chronic diseases worldwide (63). Emerging data suggest that H₂S improves diabetic endothelial dysfunction, nephropathy, retinopathy, and cardiomyopathy. Some recent studies indicate that H₂S is cytoprotective during myocardial ischemia-reperfusion injury in the setting of diabetes by alleviating oxidative stress (64). H₂S inhibits insulin secretion by activating ATP-sensitive K⁺ channels. H₂S promotes blood flow by dilating blood vessels and inhibiting platelet aggregation. H₂S has been reported to have either pro- or antiapoptotic effects on β cells. Studies on the animal models suggest that excess of H₂S in pancreatic islets may contribute to both type 1 and type 2 diabetes. H₂S has also been demonstrated to regulate insulin sensitivity. In the liver, H₂S stimulates gluconeogenesis and glycogenolysis and inhibits glucose utilization and glycogen storage. H₂S may also regulate adipose tissue lipolysis, adipokine production and inflammation; the processes important for local and systemic insulin sensitivity (65). The plasma H₂S levels are significantly reduced in type 2 diabetes patients, particularly in those with a history of cardiovascular disease. The plasma H₂S levels in patients with type 2 diabetes may reflect the HbA_{1c} levels (66).

Diabetes leads to microvascular and macrovascular complications, and diabetes-associated cognitive decline

is one of the central nervous system complications, which induces advanced brain ageing, cognitive impairment and an increased risk of dementia. Exogenous H₂S could supplement endogenous loss of H₂S and improve spatial learning and memory abilities, and the underlying mechanism identified in these studies is the Akt-GSK-3 β signalling pathway (67).

H₂S levels in patients with diabetes. H₂S reduction is related to diabetes-induced bone marrow cell (BMC) dysfunction and impaired ischemic tissue repair. Restoration of H₂S production in diabetes mice improves reparative property of BMCs (68).

H₂S is generated in pancreatic β -cells as well as in insulin target tissues including the liver, adipose tissue and skeletal muscles where it may control insulin secretion and insulin resistance. H₂S protects pancreatic β -cells against apoptosis induced oxidative stress. H₂S may act as an “intrinsic brake” against glucose-induced apoptotic death in β -cells. Exogenous H₂S has also been shown to protect β -cells from apoptosis induced by hydrogen peroxide, fatty acids, and cytokines; and through phosphorylation and activation of Akt signaling, promote cell proliferation and survival. H₂S inhibits the renin-angiotensin system in the diabetic kidney and attenuated high glucose induced mesangial cell proliferation by suppression of the MAPK signaling pathway (69).

Several studies reported that H₂S directly regulated insulin sensitivity in adipocytes. H₂S improves high glucose-induced insulin resistance. H₂S mediates the effect of L-cysteine on PIP₃ and glucose uptake. H₂S activates PI3K but inhibits PTEN. Adipogenesis participates in obesity and insulin resistance regulation. H₂S promotes adipogenesis in adipocytes. Skeletal muscle serves regulates systemic glucose metabolism. CBS and CSE are abundant in human skeletal muscles similar to their expressions in human kidney and liver. H₂S might promote glucose uptake in the muscle through sensitizing the IR-PI3K-Akt signaling pathway, thus ameliorating insulin resistance.

H₂S regulates circadian clock in the skeletal muscle. Disruption of core clock genes (CCG) contributes to the progression of metabolic disorders. H₂S synthesis enzymes (CBS, CSE and 3-MST), antioxidant genes, CCG and clock-controlled genes were all decreased in muscle of HFD-induced mice, but markers of oxidative stress and expression of oxidative stress related genes were increased. Application of H₂S donor NaHS or GSH precursor improved expression of CCG. H₂S may be an important endogenous regulator of the circadian clock and metabolic disorder (70).

Skeletal muscle is now considered to be an endocrine organ, synthesizing and secreting a variety of bioactive molecules including inflammatory cytokines, growth factors, adipokines, carnitine, and more recently

hydrogen sulfide. H₂S appears to provide protection against low oxygen and nutrient supply as well as ischemic injury in multiple organs including skeletal muscles. (71).

Diabetic macroangiopathy can cause cerebro-cardiovascular diseases and constitutes one of the major causes of death in patients with diabetes. H₂S has many beneficial roles in protecting against diabetes-induced alterations in kidney, heart and blood vessels, such as diabetic renal fibrosis, palmitic acid-induced myocardial injury and nephropathy, proliferation of VSMCs and decreased thickness of VSM. H₂S can also regulate autophagy in liver cells, H9c2 cells and colon epithelial cells. However, whether H₂S protects VSMCs in DM by regulating autophagy is unknown (72).

Mitochondria, the powerhouse of the cell, participate in multifaceted regulatory pathways in normal and pathophysiological conditions. H₂S is a regulator of the Ca²⁺ channel under different physiological conditions. It modulates Ca²⁺ concentration. H₂S inhibits Ca²⁺-induced MPTP opening in adult and old rat heart. H₂S is a strong antioxidant that protects the tissue against pathological matrix turnover, in part, by scavenging ROS. H₂S is known to inhibit the L-type Ca²⁺ channel. High Ca²⁺ activates CypD, a gatekeeper protein of MPTP, thus facilitating molecular exchange between matrix and cytoplasm causing oxidative outburst and cell death (73).

20. Hydrogen sulfide and Cardioprotection

Hydrogen sulfide, an endogenous signaling molecule, plays an important role in the physiology and pathophysiology of the cardiovascular system (74). It may be worth mentioning that the roots of the cardiovascular protective effect of H₂S can be traced back to a parallel line of studies that began in the 50s, focusing on balneotherapy (beneficial effects of H₂S-containing spring baths). The first long-acting H₂S donor, synthesized and characterized by Philip Moore's group in Singapore, was designated as GYY4137; this compound was found to exert beneficial blood pressure and vascular effects in the initial report, and was subsequently used in many dozens of follow-up studies and showed therapeutic benefit in many systems.

Hydrogen sulfide (H₂S) has been demonstrated to reduce myocardial infarction and cardiac ischaemia-reperfusion (I/R) injury. H₂S-mediated cardioprotection has been reported to involve activation of Nrf2, increased glutathione levels, and activation of endothelial nitric oxide synthase (eNOS). However, the precise details of the mechanism by which H₂S activates these pathways are poorly understood. Furthermore the observation that H₂S can elicit protective effects when administered on reperfusion suggests a role for a rapid signalling mechanism. H₂S has also been recently suggested to signal via protein S-sulfhydration (SSH), another important redox-based post-translational modification on

protein cysteine residues. H₂S signalling has also been reported to alter NO bioavailability and to alter eNOS activity. Indeed the protective effects of H₂S against I/R are lost in eNOS phosphomutant mice. Furthermore, the H₂S activation of eNOS is reported to be mediated by SSH of eNOS. Taken together, these studies suggest that H₂S-induced protection is NO signalling dependent (75).

21. Hydrogen sulfide and Cancer

In various forms of systemic inflammation, as well as several forms of cancer (76), as breast (77), gastric (78), colon (79), prostate (80) or colorectal cancer (81) - circulating H₂S levels are increased due to the upregulation of H₂S-producing enzymes. Pharmacological inhibition of H₂S biosynthesis shows therapeutic potential.

Recent studies indicate that H₂S has both pro-cancer and anti-cancer effects. Indeed, different types of cancer utilize different H₂S-associated pathways, and the final effect on cell survival or cell death appear to be dose- and tumour cell type-dependent (82).

In the field of antitumor research, endogenous H₂S induces angiogenesis, accelerates the cell cycle and inhibits apoptosis, which results in promoting oncogenesis eventually. Interestingly, high concentrations of exogenous H₂S liberated from donors suppress the growth of various tumors via inducing cellular acidification and modulating several signaling pathways involved in cell cycle regulation, proliferation, apoptosis and metastasis. The selective release of certain concentrations of H₂S from H₂S donors in the target has been considered as a tumor therapy strategy (83).

It has been shown that endogenous production of H₂S is generally low, making it difficult to elucidate the precise biological functions. Chemical compounds that could degrade in response to a specific trigger to release H₂S, termed H₂S donors, include a number of delivery systems and functional groups, some of which mimic the controlled endogenous production in response to biologically specific and relevant conditions. Hyaluronic acid (HA) is a non-sulfated glycosaminoglycan composed of two disaccharide units, D-glucuronic acid and N-acetyl-D-glucosamine, linked via β-1,3- and β-1,4-glycosidic bonds. HA has been considered one of the best biopolymers in terms of safety issues and widely used for many biomedical applications including tissue engineering and drug delivery. H₂S donors could suppress the growth of human cancer cells by inhibiting the PI3K/AKT/mTOR and RAS/RAF/MEK/ ERK signaling pathways (84). H₂S has been shown to inhibit cell survival of androgen-independent, androgen-dependent, and antiandrogen-resistant prostate cancer cells through different mechanisms. Various H₂S-releasing compounds, including sulfide salts, diallyl disulfide, diallyl trisulfide, sulfuraphane, and other polysulfides, also have been

shown to inhibit prostate cancer growth and metastasis (80).

While most studies of H₂S and cancer have focused on colonic cancer, there is considerable evidence that H₂S-based chemoprevention will be effective for a broader range of tumours, including melanoma (85).

The idea of killing cancers via intracellular acidification is not new. In fact, there are many compounds designed, and are already in clinical trials, to target carbonic anhydrases, ion exchangers or proton pumps. Yet, many are hampered by lack of selectivity for cancer cells, thence giving undesired side effects and enhanced toxicity. Unlike the inhibitors in clinical development that solely targets tumor pH regulatory protein(s), H₂S is therefore, the key component that potentiates the action of metformin and simvastatin. Of note, this combinatorial regime possesses therapeutic value, for H₂S effect has been demonstrated to be specific to cancer cells (86).

In some pathological process, as cancer and glycometabolic malfunction, different studies show that H₂S may exert different or even opposite actions. Reduced endogenous H₂S has been linked to the cause of many diseases or their stage-dependent injury. Therefore, to enhance endogenous H₂S content is of great pharmacological significance with the use of H₂S releasing/stimulating reagents. It is gratifying to see the development of H₂S releasing reagents has made great progress in recent years (87).

The fact that the mechanism of H₂S-mediated acceleration of cancer metastasis is unknown hampers the development of anti-metastasis therapies. CD36 functioned as a H₂S-targeted receptor. The use of neutralizing antibodies or inhibitors to block CD36 could accomplish an almost complete inhibition of metastasis in immunodeficient orthotopic mouse models of oral squamous cell carcinoma (88), with no side effects (89).

22. Hydrogen sulfide and Asthma

The respiratory mucosal epithelium is the first internal line of defense by acting as a major physical barrier between internal and external environments. H₂S-related enzymes are expressed in the human lungs. It is now acknowledged that H₂S is required for the development of lung vasculature and alveolarisation, and in other lung functions, including airway tone and pulmonary circulation. H₂S appears to be involved in various processes namely airway mucolytic activity, oxidative stress, inflammatory state, cell proliferation, and apoptosis. Both endogenous and exogenous H₂S show positive effects upon the respiratory tract by modulating the mucolytic activity. This appears to result from the interactions between H₂S and the disulfide bonds of mucins, resulting in breakage of the latter, which allows the mucus to become less viscous. The production of endogenous H₂S induces the opening of K⁺ATP and the

activation of the cAMP pathway. Additionally, exogenous H₂S inhibits Na⁺/K⁺-ATPase and calcium-sensitive potassium channels in human bronchiolar epithelia, thereby triggering electrolyte absorption. This leads to an increase in mucociliary clearance, and therefore the elimination of foreign microorganisms can be more effective. Inhibition of transepithelial sodium absorption (via inhibition of Na⁺/K⁺-ATPase) is favored under acute hypoxia, in order to avoid H₂S degradation, as well as exogenous H₂S exposure. Additionally, an amelioration of mucociliary function was observed with inhalation of exogenous H₂S, as confirmed by a substantial reduction of mean mucociliary transport time in patients with chronic rhinosinusitis. H₂S antioxidant features seem to involve both an indirect action and an induction of endogenous antioxidant defenses. For instance, by stimulating cysteine and cysteine transporter activity, H₂S induces an augmentation of substrate levels that are necessary for GSH production. Unlike large size antioxidants, H₂S can easily cross both plasma and mitochondrial membranes. This allows H₂S to more promptly reach its biological targets, and therefore it is considered to be more effective at diminishing cellular oxidative stresses, and at increasing antioxidant defenses. Antioxidant effects of sulphurous thermal waters (STWs), an exogenous H₂S source, provide protection against oxidative DNA damage.

The anti-inflammatory properties of H₂S can in part be explained by its potent reducing, antioxidant, and scavenging features. Nevertheless, there are controversial results concerning H₂S properties, since it appears to exert both pro- and anti-inflammatory effects. Some authors have shown an *in vitro* promotion of granulocyte survival via H₂S-induced inhibition of caspase-3-cleavage and p38 phosphorylation and in an oxidative stress environment, activated neutrophils seem to be able to convert H₂S into sulfite, which is associated with inflammation. Moreover, there are studies suggesting that this gaseous transmitter is involved in GSH depletion and ROS formation, and consequently induction of mitochondrial cell death pathways. Meanwhile, others have suggested the participation of H₂S in some key anti-inflammatory pathways, such as: (i) suppression of leukocyte adherence and migration, mediated by K⁺ATP activation in endothelial cells and leukocytes, (ii) inhibition of oxidized low-density lipoprotein-induced macrophage inflammation via NF- κ B suppression, leading to a reduction of several pro-inflammatory cytokines (e.g., IL-1 β , IL-6, and IL-8), and (iii) reduction of neutrophil toxic effects by inhibiting myeloperoxidase activity. However, these discrepancies may be related to differences in H₂S concentration. In fact, beneficial effects of H₂S generally prevail at lower concentrations, whereas deleterious effects are observed at higher levels.

Since H₂S inhibits the activity of phosphodiesterase, cGMP net levels increase. Therefore, H₂S might eventually act as a good auxiliary agent in the treatment of pulmonary hypertension (90).

23. Hydrogen sulfide and Allergic Rhinitis

Allergic Rhinitis AR is a chronic inflammatory disease that affects the nasal airways, which become inflamed and engorged after exposure to an allergen to which patients are sensitized (91). H₂S can be found in human nasal mucosa as well as in the plasma of healthy subjects. This may be partially explained by the presence of CBS and CSE in human nasal epithelium. While CBS is mainly distributed in the superficial epithelium and submucosal glands, CSE is exclusively localized in vascular endothelium and surrounding smooth muscles. In the nasal and sinus mucosa, the amount of H₂S was shown to be increased in mild and moderate/severe persistent AR. In this context, there is an enhancement of human H₂S-synthesized enzymes, mRNA and protein levels, which consequently leads to a significant increment of H₂S levels in human airway. This may indicate a compensatory mechanism to attempt to revert the pro-inflammatory state. Although there are few data specifically regarding STWs-treatment of AR, an amelioration in AR patients suffering from allergen-specific non-seasonal rhinitis when treated with a S-based compound water was observed, and this was associated with a reduction in total IgE and an increase in IgA serum levels. In accordance with this, it was suggested that STWs may exert an immunomodulatory activity by inducing an increase in IgA levels in nasal mucus. Another study also showed significant amelioration when AR patients were treated with STWs, namely in terms of a significant decrease in nasal flow resistance and nasal mucociliary transport time in 84% of subjects. Likewise, a significant reduction of IgE and an increase in IgA levels as well as an improvement of subjective symptomatology assessment scale were also observed. Data from these studies suggest a compensatory mechanism in order to reduce the presence of pro-inflammatory mediators and an improvement of the inflammatory state. However, clearly more thorough studies on the immunomodulatory and anti-inflammatory effects of STWs treatments in AR patients (as well as in patients with chronic rhinosinusitis) are needed. Due to common immunopathophysiology, there is a close relationship between AR and asthma ("single airways concept"). In fact, AR is regarded as a risk factor for the development of asthma. Thus, although few studies have been carried out with STWs treatment in patients with asthma, we shall now analyse this context (90).

24. Hydrogen sulfide and COPD

Chronic Obstructive Pulmonary Disease COPD is an airways disease caused by significant exposure to noxious particles or gases (92). H₂S metabolism may be

altered in the lung tissue of COPD patients, just as it is in smokers. H₂S levels in COPD subjects vary longitudinally, with higher levels being more frequently seen in a stable state than in acute exacerbations of COPD. Serum H₂S levels are significantly decreased in COPD subjects with very symptomatic exacerbations induced by bacteria and viruses, compared to control subjects (93). H₂S synthesis is induced in order to counter the infections-mediated exacerbations. Currently, the most frequent therapies used to treat lung viral and bacterial infections are antiviral and antibiotics. However, over time, antibiotic-resistant bacteria may be increasingly observed. To overcome these problems, some studies focused on the clinical efficacy of therapeutic S-based compounds (H₂S donors, STWs).

Despite certain contradictions founded in the previous studies, all indicated significant changes of H₂S and H₂S-synthesized enzymes levels in COPD subjects. These H₂S metabolism alterations seems to contribute, at least in part, to exacerbations and worsening of this respiratory disease state, affecting general lung function. A significant reduction of oxidative stress and an amelioration of symptoms in subjects suffering from moderate to severe COPD were observed after a 12-day inhalation with STWs and 1 month after the end of the treatment. Similar beneficial effects have been observed in patients with chronic rhinosinusitis (6). Further studies focusing on mechanisms underlying such STWs-driven improvement in COPD patients are warranted. In summary, in spite of generally effective drug-based treatments for most cases of AR, bronchial asthma, and COPD, some patients still show a sub-optimal response to such treatments. Furthermore, over time, long-term high-dose therapy may be associated with the development of some adverse effects. Thus, additional thermal spa complementary therapeutic tool mainly for subjects whose symptoms are not adequately controlled with the usual drug-based therapeutic approach seems to be a good option. With supplementary STWs treatment it may be possible to regain symptom control and eventually reduce baseline drug therapy. Another aspect which must be borne in mind is that with STWs treatment at Spas, additional psychological components may also contribute toward clinical improvement. These components include leisure time, opportunity for relaxation, being aware of regular clinical monitoring, and various cultural aspects, all of which may play a part in Spa treatment-associated final results (90).

25. Hydrogen sulfide and COVID-19

The COVID-19 pandemic caused by SARS-Cov-2 demands rapid, safe and effective therapeutic options. The "cytokine storm" syndrome, with subsequent acute lung injury (and other systemic disorders), is a frequent feature of many infectious diseases, including those caused by many viral pathogens. A cytokine storm is also

observed in the most serious cases of COVID-19. H₂S is a reducing agent and thus can be oxidized by several circulating oxidants. However, one of the main catabolic pathways for H₂S operates in the mitochondria, leading to the formation of thiosulfate. In the case of COVID-19 patients, there is a high incidence of coagulative disorders, generating a disseminated intravascular coagulation (DIC), which is lethal in severe conditions. The possibility of modulating PLA formation by H₂S is therefore of therapeutic interest, as PLAs promote both vascular inflammation and coagulation. H₂S promoted anti-inflammatory effects through epigenetic alterations. The effect of H₂S on viral infections represents a promising research field, which is presently still poorly investigated. Many pulmonary viral infections increase ROS levels and impair antioxidant enzymes, including those associated with the Nrf2–ARE pathway, with consequent reduction of antioxidant response. H₂S-releasing agents evoke significant protective effects against oxidative stress in the host cells, which are mediated by H₂S and/or related sulfane sulfur through well-clarified mechanisms, such as increased nuclear translocation of Nrf2 and inhibition of NADPH oxidase enzymes. Besides the triggering of the “antioxidant machinery” directly attributable to H₂S, other “indirect” effects are also involved in antioxidant activity of H₂S. Very recently, GSH was also proposed as a potentially useful agent against SARS-CoV-2, since high-throughput artificial intelligence-based binding affinity prediction indicated a possible GSH interaction with ACE2 or transmembrane serine protease 2 (TMPRSS2), two human proteins closely involved in the process of viral adhesion and entry into the host cell (94).

26. Hydrogen sulfide and Burns

The known role of H₂S in inflammation and sepsis underline the role of H₂S in burns. H₂S has a key role in inducing edema, which is a cardinal sign of inflammation. Understanding the role of H₂S in sepsis and shock is particularly important due to the high mortality associated with both conditions. From current studies, we can postulate that H₂S would act as a pro-inflammatory agent in an early burn via a substance P-mediated increase in permeability and edema. H₂S seems to have a conflicting role in sepsis, and the underlying mechanism needs to be studied in various animal models at different intervals of sepsis development. H₂S plays a novel role in burns, and it appears that the modulation of H₂S administration may have a therapeutic role. However, the role of H₂S is complex, and the effect of H₂S on the inflammatory response varies at different time points following the injury. Studies need to focus on delineating the pathways and mechanisms by which H₂S induces and inhibits inflammation. The optimal concentration at which H₂S inhibits inflammation also needs to be determined. Studying the role of H₂S at each

stage will enable us to capture a complete picture of the exact role of H₂S following a burn. H₂S appears to have two very different roles: as a mediator of inflammation and as an inhibitor of inflammation. Severe burns are characterized by an inflammatory response that can progress to a systemic response and lead to sepsis, shock and multiorgan failure. Therefore, identifying agents that can reduce the duration and complications of burns is of great interest. It appears that H₂S acts to promote inflammation in the initial period post-injury, but at later stages, H₂S reduces inflammation and improves wound healing. This biphasic action of H₂S in burns is a novel concept, and although this action has not yet been extensively studied, it is promising due to the potential therapeutic role of H₂S. The development of H₂S as a therapeutic agent is currently still in the early stages; however, the potential of H₂S for treating burns is significant (18).

27. Hydrogen sulfide and Skin diseases

Although sulfur-rich thermal waters have ancestrally been used in the context of dermatological conditions (95), a global mapping of the molecular effects exerted by H₂S on human keratinocytes is still lacking. Most of the scientific literature on H₂S effects on cell viability, proliferation, activation, cytokine secretion and adhesion. As far as skin is concerned, sulfurs are able to penetrate the skin and a sulfur-rich balneotherapy (96), known to be effective in the treatment of psoriasis (97), may be useful also in other T-cell-mediated autoimmune diseases of the skin. Sulfur has a long history of topical use: hydrogen sulfide is present in a variety of skin care products, including soaps and cleansers, and is used to reduce dandruff and decrease scalp scaling and dryness. It has recognized fungicidal, bactericidal and antipruritic properties, and therefore it is used for the therapy of mite infestations in animals. Sulfur was even proposed as an antidote for acute exposure to radioactive material. Indeed, it has been demonstrated that sulfurs favor wound healing, acts as a keratolytic agent and can induce various histological modifications in the skin (including hyperkeratosis, acanthosis, and dilation of dermal vessels) (98).

Vascular endothelial growth factor (VEGF) is an important mediator involved in inflammatory processes associated with aberrant angiogenesis. In skin inflammation, epidermal cells are important VEGF-producing cells which probably contribute to the pathogenesis of diverse epidermal diseases through their chemotactic activity on inflammatory cells and through their angiogenic effect. H₂S significantly increases NO, in a manner Akt-dependent, in keratinocyte cells. H₂S as both NO donor and anti-VEGF, may be a promising therapeutic for chronic inflammatory disorders of the skin, i.e. psoriasis (99).

A recent study showed that 10-100 μM NaHS and the overexpression of CSE could promote cell proliferation and melanin synthesis by increasing the production of melanogenic enzymes in primary human epidermal melanocytes. Therefore, pharmacologic regulation of H₂S may be a novel strategy for skin disorders caused by the loss of melanocytes and dysregulation of oxidative stress. Abnormal H₂S metabolism is associated with the pathogenesis of fibrosis, causing damage to the structure and function of tissues and organs. Several *in vivo* and *in vitro* studies have shown that both endogenous H₂S levels and the expression of H₂S-related enzymes in plasma are significantly decreased in fibrotic diseases, but supplementation with exogenous H₂S could alleviate the severity of fibrosis. H₂S could restore a normal morphologic phenotype in Werner syndrome fibroblasts by attenuating oxidative damage and modulating the mTOR pathway. Beneficial effects of H₂S were demonstrated on systemic sclerosis associated skin and lung fibrosis (100).

H₂S contained in sulfur-rich spa waters is a bio-active component able to modify the immunological status of non-diseased human keratinocytes (101). Further studies are needed to establish causal links between the above-identified biological effects and the previously reported medical benefits afforded by sulfur-rich spa waters in conditions such as psoriasis, eczema or wound healing (102).

28. H₂S role on Gynecological diseases

Since H₂S has a tocolytic action, is produced from all reproductive tissues studied so far and the fact that they all express CBS and CSE, it does seem that H₂S has a functional role in the female reproductive system.

For example, H₂S could be involved in maintaining uterine quiescence during pregnancy, and there are some parallels with NO in this action. It is possible that upregulation of CBS and CSE in pregnancy causes local production of H₂S in the myometrium and the placenta, which would inhibit uterine contractions. In rats it has been shown that externally administered H₂S prolonged the duration of labor. H₂S donors have been shown to inhibit spontaneous uterine contractions in rats *in vitro*. So far, there are no data on the effects of H₂S on human myometrium. It is now well established that H₂S can act as a vasodilator and is involved in the regulation of blood pressure, at least in the mouse and rat. However, H₂S can also cause vasoconstriction. These apparently contradictory effects of H₂S appear to be related to its concentration; generally H₂S causes vasoconstriction at low concentrations and vasodilation at high concentrations. H₂S could be involved in the local control of blood flow, particularly in the placenta; however, there are currently no published studies on human placental blood vessels. We have shown that H₂S production from human placental homogenates was

significantly greater in a low-oxygen environment compared with a normal-oxygen environment, suggesting that under hypoxic conditions *in vivo*, increased H₂S release may cause local vasodilation. The CO/heme oxygenase system has been shown to function in a similar way. As pre-eclampsia is thought to involve placental ischemia, it does raise the fascinating possibility of whether a lack of H₂S production locally would cause a failure of reflex vasodilation. It has been suggested that the H₂S system may act as a back up to the NO system in vasodilation, with failure of both systems leading to disease. Olson et al. have proposed that H₂S is part of an oxygen sensing system in the vasoconstriction and vasodilation response to hypoxia. It is thought that the rate of metabolism of H₂S serves as the oxygen sensor in vascular smooth muscle. In Olson's proposed model, the concentration of vasoactive H₂S is regulated by the balance between vascular H₂S production and its oxidation by available oxygen. Clearly under hypoxic conditions the lack of oxygen would decrease the oxidation of H₂S and its vascular effects would persist. Perhaps competition between oxygen and H₂S for binding to cytochrome oxidase in mitochondria could also be involved. If this model is correct then there could be implications for pre-eclampsia.

CBS appears to be essential for female reproductive function, as it has been shown that *CBS*^{-/-} knockout mice are infertile (103).

Endometriosis, one of the most frequently encountered gynecological disorders affecting up to 10% of the reproductive-aged female population, is characterized by the presence and growth of endometrial tissue outside the uterine cavity. Ectopic endometrium expresses a high level of CBS and CSE. Meanwhile, we identified that exogenous and endogenous H₂S could induce cell proliferation, and this pro-proliferation effect was mediated by activation of NF- κ B signaling (104).

29. Hydrogen sulfide and Clinical trials

Postoperative pain relief (105) facilitates recovery from injury, but optimal perioperative or postoperative treatments remain problematic. In recent years, the biological importance of H₂S in many organs and systems has become increasingly clear. Extensive studies have documented the many beneficial antiinflammatory effects of this gaseous mediator. Several groups have developed novel H₂S-releasing drugs in the hope that they will be useful for treatment of a range of diseases.

There is a large body of evidence regarding the ability of H₂S to reduce inflammation and pain (106). In the Gastro Intestinal (GI) tract, H₂S has been shown to reduce inflammation and accelerate healing of damaged tissue (such as ulcers), while suppression of H₂S production in the GI tract results in impaired healing of tissue injury and exacerbation of inflammation (107).

The plasma H₂S levels were measured in 154 outpatients with type 2 diabetes and 66 outpatients without diabetes (66). The plasma H₂S levels were measured using the methylene blue assay. The patients with type 2 diabetes showed a progressive reduction in the plasma H₂S levels (45.1±15.5 μM versus 54.0±26.4 μM, p<0.05), which paralleled poor glycemic control. There was a significant correlation between a reduction in the plasma H₂S levels and the HbA_{1c} levels (β=-0.505, p<0.01). Furthermore, a reduction in the plasma H₂S levels was found to be related to a history of cardiovascular diseases in patients with type 2 diabetes (39.9±13.8 μM versus 47.5±15.9 μM, p<0.01) (66).

A proof-of-concept, Phase 2 clinical trial of the gastrointestinal safety of a hydrogen sulfide-releasing anti-inflammatory drug (107): 244 healthy volunteers completed a 2-week, double-blind study, taking either ATB-346 (250 mg once daily) or naproxen (550 mg twice daily), with upper GI ulceration being examined endoscopically. ATB-346 is an H₂S-releasing derivative of naproxen. Naproxen is among the most widely used NSAID for treatment of arthritis and other conditions characterized by inflammation and pain. In 2016, Antibe performed a Phase 2 clinical trial, in osteoarthritis patients, to test the hypothesis that ATB-346 was more potent and long acting than naproxen. The study involved 12 osteoarthritis patients taking ATB-346 (250 mg) once daily for 10 days, with measurements of the patients' level of pain during the 10-day period. Once daily administration of ATB-346 at a dose of 250 mg could produce substantial relief of pain in osteoarthritis patients (108).

Of the 126 subjects treated with naproxen, 53 had at least one ulcer greater than 3 mm in diameter. Of the 118 subjects treated with ATB-346, three had at least one ulcer greater than 3 mm in diameter. The incidence of ulcers between the two groups was significantly different (P < 0.05). There were also far more gastric ulcers (50-fold) in the naproxen group than in the ATB-346 group. There was a total of 210 gastric ulcers observed in the 53 subjects that developed ulcers, while in the ATB-346 group, there was a total of only four gastric ulcers in the three subjects that developed ulcers. Ulceration of the duodenal mucosa was also observed in seven of the subjects treated with naproxen (with an average of 3.4 duodenal ulcers per subject), while duodenal ulcers were not observed in any of the subjects treated with ATB-346.

Plasma levels of H₂S levels in subjects treated with ATB-346 were significantly greater (P < 0.05) than those in subjects treated with naproxen. Plasma H₂S levels were measured by a modified MBB method. Hence, this biomolecule-bound sulfide pool serves as a sulfide buffer system by responding to concentration changes via shifting chemical equilibria. Importantly, when sulfide is

consumed in a biological process, the sulfide pool will also likely serve as a slow endogenous sulfide donor system and maintain steady-state sulfide levels via similar mechanisms. Therefore, the differences measured by these methods are highly relevant and represent the sum of differences in (a) free sulfide levels and (b) the ability of the sulfide pool to release sulfide.

Cumulative data from the three clinical trials in which ATB-346 has been administered at 250 mg once daily for 10–14 days reveal a 4.7% overall incidence of clinically significant increases in liver transaminases. This is comparable to what has been observed with the use of NSAIDs such as diclofenac, naproxen, and piroxicam (107).

30. Conclusions

Hydrogen sulfide (H₂S) is a signaling molecule that is actively synthesized in the tissues and is involved in the regulation of vascular tone, neuromodulation, cytoprotection, inflammation and apoptosis. In recent years, new data on animal and human H₂S metabolism and function under the effect of various endogenous and exogenous factors, including drugs were collected. H₂S metabolism and regulation, peculiarities of transport, signaling, biological role and participation in pathogenesis are focus topics of research. It was demonstrated that vitamin-microelement and microelement complexes with H₂S-synthesizing enzymes cofactors and activators represent a promising approach for H₂S content correction in tissues. Although the field of H₂S biology has dramatically expanded over the last decade, many topics and issues remain that will demand continuing attention. Mainly due the presence of H₂S, STWs might be an advantageous and promising option as an add-on non-pharmacological complementary therapy for relatively long-lasting improvement of clinical parameters in patients with reviewed diseases. Furthermore, H₂S-rich STWs have various effects upon inflammatory and immunological parameters that may contribute toward their clinical efficacy.

31. Funding

This research did not receive any specific grant from funding public, commercial, or not-for-profit agencies.

32. Author contribution

All authors have consistently contributed to this article, CM is the lead author who suggested the idea of the article. CM and DM participated in the process of systematic review, writing and verification of the article. GO coordinated and advised the progress of the article.

33. Declaration of interests

This article does not contain any studies with human or animal subjects. This study did not require written consent from patients. The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References:

1. Tomasova L, Konopelski P, Ufnal M. Gut bacteria and hydrogen sulfide: The new old players in circulatory system homeostasis. *Molecules*. 2016;21(11):1–18.
2. Aroca A, Gotor C, Bassham DC, Romero LC. Hydrogen sulfide: From a toxic molecule to a key molecule of cell life. *Antioxidants*. 2020;9(7):1–24.
3. Munteanu C, Dogaru G, Rotariu M, Onose G. Therapeutic gases used in balneotherapy and rehabilitation medicine - scientific relevance in the last ten years (2011 – 2020) - Synthetic literature review. 2021;111–22.
4. Deja-Sikora E, Gołębiewski M, Kalwasińska A, Krawiec A, Kosobucki P, Walczak M. Comamonadaceae OTU as a Remnant of an Ancient Microbial Community in Sulfidic Waters. *Microb Ecol*. 2019;78(1):85–101.
5. Gris B, Treu L, Zampieri RM, Caldara F, Romualdi C, Campanaro S, et al. Microbiota of the therapeutic Euganean thermal muds with a focus on the main cyanobacteria species. *Microorganisms*. 2020;8(10):1–23.
6. Hale VL, Jeraldo P, Mundy M, Yao J, Keeney G, Scott N, et al. Synthesis of multi-omic data and community metabolic models reveals insights into the role of hydrogen sulfide in colon cancer. 2018; Available from: <https://doi.org/10.1016/j.ymeth.2018.04.024>
7. Cao X, Ding L, Xie ZZ, Yang Y, Whiteman M, Moore PK, et al. A Review of Hydrogen Sulfide Synthesis, Metabolism, and Measurement: Is Modulation of Hydrogen Sulfide a Novel Therapeutic for Cancer? *Antioxidants Redox Signal*. 2019;31(1).
8. Panthi S, Manandhar S, Gautam K. Hydrogen sulfide, nitric oxide, and neurodegenerative disorders. *Transl Neurodegener*. 2018;7(1):1–8.
9. Malagrino F, Zuhra K, Mascolo L, Mastronicola D, Vicente JB, Forte E, et al. Hydrogen sulfide oxidation: Adaptive changes in mitochondria of SW480 colorectal cancer cells upon exposure to hypoxia. *Oxid Med Cell Longev*. 2019;2019.
10. Szabo C. Hydrogen Sulfide, an Endogenous Stimulator of Mitochondrial Function in Cancer Cells. *Cells*. 2021;10(2).
11. Gambari L, Grigolo B, Grassi F. Hydrogen sulfide in bone tissue regeneration and repair: State of the art and new perspectives. *Int J Mol Sci*. 2019;20(20).
12. Salloom FN. Hydrogen sulfide and cardioprotection- Mechanistic insights and clinical translatability. 2015; Available from: <http://dx.doi.org/10.1016/j.pharmthera.2015.04.004>
13. Xiao Q, Ying J, Xiang L, Zhang C. The biologic effect of hydrogen sulfide and its function in various diseases. *Med (United States)*. 2018;97(44).
14. Szabo C. A timeline of hydrogen sulfide (H₂S) research: From environmental toxin to biological mediator. 2017; Available from: <https://doi.org/10.1016/j.bcp.2017.09.010>
15. Wator K, Dobrzyński D, Sugimori K, Kmiecik E. Redox potential research in the field of balneotherapy: case study on equilibrium approach to bioactive elements in therapeutic waters. *Int J Biometeorol*. 2020;64(5):815–26.
16. Wang R. Physiological implications of hydrogen sulfide: A whiff exploration that blossomed. *Physiol Rev*. 2012;92(2):791–896.
17. Cao X, Cao L, Ding L, Bian J song. A New Hope for a Devastating Disease: Hydrogen Sulfide in Parkinson's Disease. *Mol Neurobiol*. 2018;55(5):3789–99.
18. Akter F. The role of hydrogen sulfide in burns. *Burns* [Internet]. 2016;42(3):519–25. Available from: <http://dx.doi.org/10.1016/j.burns.2015.07.005>
19. Stein A, Bailey SM. Redox biology of hydrogen sulfide: Implications for physiology, pathophysiology, and pharmacology. *Redox Biol* [Internet]. 2013;1(1):32–9. Available from: <http://dx.doi.org/10.1016/j.redox.2012.11.006>
20. Liu J, Duan C, Zhang W, Ta HT, Yuan J, Zhang R, et al. Responsive nanosensor for ratiometric luminescence detection of hydrogen sulfide in inflammatory cancer cells. *Anal Chim Acta*. 2020;1103:156–63.
21. Song X, Dong B, Kong X, Wang C, Zhang N, Lin W. A cancer cell-specific two-photon fluorescent probe for imaging hydrogen sulfide in living cells. *RSC Adv*. 2017;7(26):15817–22.
22. Bilska-Wilkosz A, Iciek M, Kowalczyk-Pachel D, Górny M, Sokolowska-Jezewicz M, Wlodek L. Lipoic acid as a possible pharmacological source of hydrogen sulfide/sulfane sulfur. *Molecules*. 2017;22(3).
23. Rapolienė L, Razbadauskas A, Jurgelėnas A. The Reduction of Distress Using Therapeutic Geothermal Water Procedures in a Randomized Controlled Clinical Trial. *Adv Prev Med*. 2015;2015:1–10.
24. Barnich N, Rodrigues M, Sauvanet P, Chevarin C, Denis S, Le Goff O, et al. Beneficial effects of natural mineral waters on intestinal inflammation and the mucosa-associated microbiota. *Int J Mol Sci*. 2021;22(9):1–14.
25. Munteanu C, Rotariu M, Dogaru G, Ionescu VE, Vlad C, Onose G. Mud therapy and rehabilitation - scientific relevance in the last six years (2015 – 2020) Systematic literature review and meta-analysis based on the PRISMA paradigm. 2021;12(1):1–15.
26. Altaany Z, Alkaraki A, Abu-siniyeh A, Al Momani W, Taani O. Evaluation of antioxidant status and oxidative stress markers in thermal sulfurous springs residents. *Heliyon* [Internet]. 2019;5(11):e02885. Available from: <https://doi.org/10.1016/j.heliyon.2019.e02885>
27. Cheleschi S, Gallo I, Tenti S. A comprehensive analysis to understand the mechanism of action of balneotherapy: why, how, and where they can be used? Evidence from in vitro studies performed on human and animal samples. *Int J Biometeorol*. 2020;64(7):1247–61.
28. Burguera EF, Vela-Anero Á, Gato-Calvo L, Vaamonde-García C, Meijide-Faílde R, Blanco FJ. Hydrogen sulfide biosynthesis is impaired in the osteoarthritic joint. *Int J Biometeorol*. 2020;64(6):997–1010.
29. Fioravanti A, Bender T, Karagülle Z. Balneotherapy in osteoarthritis: Facts, fiction and gaps in knowledge. 2017; Available from: <http://dx.doi.org/10.1016/j.eujim.2017.01.001>
30. Silva A, Oliveira AS, Vaz C V., Correia S, Ferreira R, Breitenfeld L, et al. Anti-inflammatory potential of Portuguese thermal waters. *Sci Rep* [Internet]. 2020;10(1):1–13. Available from: <https://doi.org/10.1038/s41598-020-79394-9>
31. Pavlovska I, Klavina A, Auce A, Vanadzins I, Silova A, Komarovska L, et al. Assessment of sapropel use for pharmaceutical products according to legislation,

- pollution parameters, and concentration of biologically active substances. *Sci Rep* [Internet]. 2020;10(1):1–12. Available from: <https://doi.org/10.1038/s41598-020-78498-6>
32. Fonseca LJS Da, Nunes-Souza V, Goulart MOF, Rabelo LA. Oxidative Stress in Rheumatoid Arthritis: What the Future Might Hold regarding Novel Biomarkers and Add-On Therapies. *Oxid Med Cell Longev*. 2019;2019.
 33. Kuciel-Lewandowska J, Kasperczak M, Paprocka-Borowicz M. Assessment of Changes in the Hemoglobin Level under the Influence of Comprehensive Spa Therapy Using Therapeutic Radon-Sulfur Waters and Its Correlation with Free Radical Reactions. *Evidence-based Complement Altern Med*. 2020;2020.
 34. Gambari L, Grigolo B, Filardo G, Grassi F. Sulfurous thermal waters stimulate the osteogenic differentiation of human mesenchymal stromal cells – An in vitro study. *Biomed Pharmacother* [Internet]. 2020;129(June):110344. Available from: <https://doi.org/10.1016/j.biopha.2020.110344>
 35. Sieghart D, Liszt M, Wanivenhaus A, Bröll H, Kiener H, Klösch B, et al. Hydrogen sulphide decreases IL-1 β -induced activation of fibroblast-like synoviocytes from patients with osteoarthritis. *J Cell Mol Med*. 2015;19(1):187–97.
 36. Vela-Anero A, Hermida-G Omez T, Gato-Calvo L, Vaamonde-García C, Díaz-Prado S, Meijide-Faílde R, et al. Long-term effects of hydrogen sulfide on the anabolic-catabolic balance of articular cartilage in vitro. 2017; Available from: <http://dx.doi.org/10.1016/j.niox.2017.08.004>
 37. Lv X, Wang J, Bao Y, Tang Y, Xing W, Wu Q, et al. The effectiveness of balneotherapy and aquatic exercise on bone metabolism: a systematic review and meta-analysis. *Complement Ther Clin Pract* [Internet]. 2021;101429. Available from: <https://doi.org/10.1016/j.ctcp.2021.101429>
 38. Boros M, Kemény Á, Seb" Ok D , Teréz Bagoly B, Perkecz A, Pet" Oházi Z, Maász G, et al. Sulphurous medicinal waters increase somatostatin release: It is a possible mechanism of anti-inflammatory effect of balneotherapy in psoriasis. *Eur J Integr Med* [Internet]. 2013;5:109–18. Available from: www.sciencedirect.com
 39. Yang X, Wang C, Zhang X, Chen S, Chen L, Lu S, et al. Redox regulation in hydrogen sulfide action: From neurotoxicity to neuroprotection. *Neurochem Int*. 2019;128(March):58–69.
 40. Shan H, Qiu J, Chang P, Chu Y, Gao C, Wang H, et al. Exogenous Hydrogen Sulfide Offers Neuroprotection on Intracerebral Hemorrhage Injury Through Modulating Endogenous H₂S Metabolism in Mice. *Front Cell Neurosci*. 2019;13(August):1–14.
 41. Zhu Y, Shui M, Liu X, Hu W, Wang Y. Increased autophagic degradation contributes to the neuroprotection of hydrogen sulfide against cerebral ischemia/reperfusion injury. *Metab Brain Dis*. 2017;32(5):1449–58.
 42. Dou Y, Wang Z, Chen G. The role of hydrogen sulfide in stroke. *Med Gas Res*. 2016;6(2):79–84.
 43. Ji J, Xiang P, Li T, Lan L, Xu X, Lu G, et al. NOSH-NBP, a novel nitric oxide and hydrogen Sulfide-Releasing hybrid, attenuates ischemic Stroke-Induced neuroinflammatory injury by modulating microglia polarization. *Front Cell Neurosci*. 2017;11(May):1–19.
 44. Liu H, Wang Y, Xiao Y, Hua Z, Cheng J, Jia J. Hydrogen Sulfide Attenuates Tissue Plasminogen Activator-Induced Cerebral Hemorrhage Following Experimental Stroke. *Transl Stroke Res* [Internet]. 2016;7(3):209–19. Available from: <http://dx.doi.org/10.1007/s12975-016-0459-5>
 45. Narne P, Pandey V, Phanithi PB. Role of Nitric Oxide and Hydrogen Sulfide in Ischemic Stroke and the Emergent Epigenetic Underpinnings. *Mol Neurobiol*. 2019;56(3):1749–69.
 46. Juman S, Nara Y, Yasui N, Negishi H, Okuda H, Takado N, et al. Reduced production of hydrogen sulfide and sulfane sulfur due to low cystathionine β -synthase levels in brain astrocytes of stroke-prone spontaneously hypertensive rats. *Biol Pharm Bull*. 2016;39(12):1932–8.
 47. Wu YH, Du HP, Xu Y, Xu ZH. Long non-coding ribonucleic acid small nucleolar host gene 1 regulates the ischemic stroke by targeting micro ribonucleic acid-376a-mediated cystathionine β synthase/hydrogen sulfide axis pathway. *Indian J Pharm Sci*. 2021;82(7):90–7.
 48. Liu Y, Liao S, Quan H, Lin Y, Li J, Yang Q. Involvement of microRNA-135a-5p in the Protective Effects of Hydrogen Sulfide Against Parkinson's Disease. *Cell Physiol Biochem*. 2016;40(1–2):18–26.
 49. Kida K, Yamada M, Tokuda K, Marutani E, Kakinohana M, Kaneki M, et al. Inhaled hydrogen sulfide prevents neurodegeneration and movement disorder in a mouse model of Parkinson's disease. *Antioxidants Redox Signal*. 2011;15(2):343–52.
 50. Hu LF, Lu M, Tiong CX, Dawe GS, Hu G, Bian JS. Neuroprotective effects of hydrogen sulfide on Parkinson's disease rat models. *Aging Cell*. 2010;9(2):135–46.
 51. Lee M, Tazzari V, Giustarini D, Rossi R, Sparatore A, Del Soldato P, et al. Effects of hydrogen sulfide-releasing L-DOPA derivatives on glial activation: Potential for treating Parkinson disease. *J Biol Chem* [Internet]. 2010;285(23):17318–28. Available from: <http://dx.doi.org/10.1074/jbc.M110.115261>
 52. Xie L, Hu LF, Teo XQ, Tiong CX, Tazzari V, Sparatore A, et al. Therapeutic Effect of Hydrogen Sulfide-Releasing L-Dopa Derivative ACS84 on 6-OHDA-Induced Parkinson's Disease Rat Model. *PLoS One*. 2013;8(4).
 53. Wang M, Tang JJ, Wang LX, Yu J, Zhang L, Qiao C. Hydrogen sulfide enhances adult neurogenesis in a mouse model of Parkinson's disease. *Neural Regen Res*. 2021;16(7):1353–8.
 54. Zhao FL, Fang F, Qiao PF, Yan N, Gao D, Yan Y. AP39, a Mitochondria-Targeted Hydrogen Sulfide Donor, Supports Cellular Bioenergetics and Protects against Alzheimer's Disease by Preserving Mitochondrial Function in APP/PS1 Mice and Neurons. *Oxid Med Cell Longev*. 2016;2016.
 55. Giovino D, Bursac B, Sbodio JI, Nalluru S, Vignane T, Snowman AM, et al. Hydrogen sulfide is neuroprotective in Alzheimer's disease by sulphydrating GSK3 β and inhibiting Tau hyperphosphorylation. *Proc Natl Acad Sci* [Internet]. 2021 Jan 26 [cited 2021 Jun 20];118(4). Available from:

- <https://www.pnas.org/content/118/4/e2017225118>
56. Kamat PK, Kyles P, Kalani A, Tyagi N. Hydrogen Sulfide Ameliorates Homocysteine-Induced Alzheimer's Disease-Like Pathology, Blood-Brain Barrier Disruption, and Synaptic Disorder. *Mol Neurobiol*. 2016;53(4):2451–67.
 57. Vandini E, Ottani A, Zaffe D, Calevro A, Canalini F, Cavallini GM, et al. Mechanisms of hydrogen sulfide against the progression of severe Alzheimer's disease in transgenic mice at different ages. *Pharmacology*. 2019;103(1–2):93–100.
 58. Paul BD, Snyder SH. Impaired redox signaling in Huntington's disease: Therapeutic implications. *Front Mol Neurosci*. 2019;12(March):1–13.
 59. Liu C, Xu X, Gao J, Zhang T, Yang Z. Hydrogen Sulfide Prevents Synaptic Plasticity from VD-Induced Damage via Akt/GSK-3 β Pathway and Notch Signaling Pathway in Rats. *Mol Neurobiol* [Internet]. 2016;53(6):4159–72. Available from: <http://dx.doi.org/10.1007/s12035-015-9324-x>
 60. Masiero S, Maccarone MC, Magro G. Balneotherapy and human immune function in the era of COVID-19. *Int J Biometeorol*. 2020;64(8):1433–4.
 61. Gálvez I, Torres-Piles S, Ortega-Rincón E. Balneotherapy, immune system, and stress response: A hormetic strategy? *Int J Mol Sci*. 2018;19(6).
 62. Dilek N, Papapetropoulos A, Toliver-Kinsky T, Szabo C. Hydrogen sulfide: An endogenous regulator of the immune system. *Pharmacol Res* [Internet]. 2020;161(June):105119. Available from: <https://doi.org/10.1016/j.phrs.2020.105119>
 63. Durante W. Hydrogen sulfide therapy in diabetes-accelerated atherosclerosis: A whiff of success. *Diabetes*. 2016;65(10):2832–4.
 64. Xie L, Gu Y, Wen M, Zhao S, Wang W, Ma Y, et al. Hydrogen sulfide induces Keap1 S-sulfhydration and suppresses diabetes-accelerated atherosclerosis via Nrf2 activation. *Diabetes*. 2016;65(10):3171–84.
 65. Bełtowski J, Wójcicka G, Jamroz-Wisniewska A. Hydrogen sulfide in the regulation of insulin secretion and insulin sensitivity: Implications for the pathogenesis and treatment of diabetes mellitus. Vol. 149, *Biochemical Pharmacology*. Elsevier Inc.; 2018. p. 60–76.
 66. Suzuki K, Sagara M, Aoki C, Tanaka S, Aso Y. Clinical implication of plasma hydrogen sulfide levels in Japanese patients with type 2 diabetes. *Intern Med*. 2017;56(1):17–21.
 67. Ma S, Zhong D, Ma P, Li G, Hua W, Sun Y, et al. Exogenous hydrogen sulfide ameliorates diabetes-associated cognitive decline by regulating the mitochondria-mediated apoptotic pathway and IL-23/IL-17 expression in db/db mice. *Cell Physiol Biochem*. 2017;41(5):1838–50.
 68. Cheng Z, Shen X, Jiang X, Shan H, Cimini M, Fang P, et al. Hyperhomocysteinemia potentiates diabetes-impaired EDHF-induced vascular relaxation: Role of insufficient hydrogen sulfide. *Redox Biol*. 2018;16(Feb.):215–25.
 69. Gheibi S, Samsonov AP, Gheibi S, Vazquez AB, Centre D, Sciences B, et al. hydrogen sulfide: Implications in diabetes. 2021;1–46.
 70. Zhang H, Huang Y, Chen S, Tang C, Wang G, Du J, et al. Hydrogen sulfide regulates insulin secretion and insulin resistance in diabetes mellitus, a new promising target for diabetes mellitus treatment? A review. *J Adv Res* [Internet]. 2021;27:19–30. Available from: <https://doi.org/10.1016/j.jare.2020.02.013>
 71. Bitar MS, Nader J, Al-Ali W, Al Madhoun A, Arefanian H, Al-Mulla F. Hydrogen sulfide donor NaHS improves metabolism and reduces muscle atrophy in type 2 diabetes: Implication for understanding sarcopenic pathophysiology. *Oxid Med Cell Longev*. 2018;2018.
 72. Qiu X, Liu K, Xiao L, Jin S, Dong J, Teng X, et al. Alpha-lipoic acid regulates the autophagy of vascular smooth muscle cells in diabetes by elevating hydrogen sulfide level. *Biochim Biophys Acta - Mol Basis Dis* [Internet]. 2018;1864(11):3723–38. Available from: <https://doi.org/10.1016/j.bbadis.2018.09.005>
 73. Sashi Papu John A, Kundu S, Pushpakumar S, Amin M, Tyagi SC, Sen U. Hydrogen sulfide inhibits Ca²⁺-induced mitochondrial permeability transition pore opening in type-1 diabetes. *Am J Physiol - Endocrinol Metab*. 2019;317(2):E269–83.
 74. Zhang Y, Li H, Zhao G, Sun A, Zong NC, Li Z, et al. Hydrogen Sulfide Attenuates the Recruitment of CD11b + Gr-1 + Myeloid Cells and Regulates Bax/Bcl-2 Signaling in Myocardial Ischemia Injury. *Sci Rep*. 2014;4:1–7.
 75. Sun J, Aponte AM, Menazza S, Gucek M, Steenbergen C, Murphy E. Additive cardioprotection by pharmacological postconditioning with Hydrogen sulfide and nitric oxide donors in mouse heart: S-sulfhydration vs. S-nitrosylation. *Cardiovasc Res*. 2016;110(1):96–106.
 76. Chen S, Yue T, Huang Z, Zhu J, Bu D, Wang X, et al. Inhibition of hydrogen sulfide synthesis reverses acquired resistance to 5-FU through miR-215-5p-EREG/TYMS axis in colon cancer cells. *Cancer Lett*. 2019;466(September):49–60.
 77. Youness RA, Gad AZ, Sanber K, Ahn YJ, Lee GJ, Khallaf E, et al. Targeting hydrogen sulphide signaling in breast cancer. *J Adv Res* [Internet]. 2021;27:177–90. Available from: <https://doi.org/10.1016/j.jare.2020.07.006>
 78. Sekiguchi F, Sekimoto T, Ogura A, Kawabata A. Endogenous Hydrogen Sulfide Enhances Cell Proliferation of Human Gastric Cancer AGS Cells. *Biol Pharm Bull*. 2016;39(5):887–90.
 79. Oláh G, Módis K, Törö G, Hellmich MR, Szczesny B, Szabo C. Role of endogenous and exogenous nitric oxide, carbon monoxide and hydrogen sulfide in HCT116 colon cancer cell proliferation. *Biochem Pharmacol*. 2018;149:186–204.
 80. Liu M, Wu L, Montaut S, Yang G. Hydrogen sulfide signaling axis as a target for prostate cancer therapeutics. *Prostate Cancer*. 2016;2016.
 81. Sakuma S, Minamino S, Takase M, Ishiyama Y, Hosokura H, Kohda T, et al. Hydrogen sulfide donor GYY4137 suppresses proliferation of human colorectal cancer Caco-2 cells by inducing both cell cycle arrest and cell death. *Heliyon* [Internet]. 2019;5(8):e02244. Available from: <https://doi.org/10.1016/j.heliyon.2019.e02244>
 82. De Cicco P, Ercolano G, Rubino V, Terrazzano G, Ruggiero G, Cirino G, et al. Modulation of the functions of myeloid-derived suppressor cells: a new strategy of

- hydrogen sulfide anti-cancer effects. *Br J Pharmacol*. 2020;177(4):884–97.
83. Li H, Xu F, Gao G, Gao X, Wu B, Zheng C, et al. Hydrogen sulfide and its donors: Novel antitumor and antimetastatic therapies for triple-negative breast cancer. *Redox Biol*. 2020;34(March):0–4.
 84. Dong Q, Yang B, Han J-G, Zhang M-M, Liu W, Zhang X, et al. Original Articles A novel hydrogen sulfide-releasing donor, HA-ADT, suppresses the growth of human breast cancer cells through inhibiting the PI3K/AKT/mTOR and Ras/Raf/MEK/ERK signaling pathways. 2019; Available from: <https://doi.org/10.1016/j.canlet.2019.04.031>
 85. Ianaro A, Cirino G, Wallace JL. Hydrogen sulfide-releasing anti-inflammatory drugs for chemoprevention and treatment of cancer. *Pharmacol Res* [Internet]. 2016;111:652–8. Available from: <http://dx.doi.org/10.1016/j.phrs.2016.07.041>
 86. Lee ZW, Zhou J, Chen CS, Zhao Y, Tan CH, Li L, et al. The slow-releasing Hydrogen Sulfide donor, GYY4137, exhibits novel anti-cancer effects in vitro and in vivo. *PLoS One*. 2011;6(6):5–11.
 87. Yang CT, Chen L, Xu S, Day JJ, Li X, Xian M. Recent development of hydrogen sulfide releasing/stimulating reagents and their potential applications in cancer and glycometabolic disorders. *Front Pharmacol*. 2017;8(SEP).
 88. Zhang S, Bian H, Li X, Wu H, Bi Q, Yan Y, et al. Hydrogen sulfide promotes cell proliferation of oral cancer through activation of the COX2/AKT/ERK1/2 axis. *Oncol Rep*. 2016;35(5):2825–32.
 89. Wang R, Tao B, Fan Q, Wang S, Chen L, Zhang J, et al. Fatty-acid receptor CD36 functions as a hydrogen sulfide-targeted receptor with its Cys333–Cys272 disulfide bond serving as a specific molecular switch to accelerate gastric cancer metastasis. *EBioMedicine* [Internet]. 2019;45:108–23. <https://doi.org/10.1016/j.ebiom.2019.06.037>
 90. Viegas J, Esteves AF, Cardoso EM, Arosa FA, Vitale M, Taborda-Barata L. Biological effects of thermal water-associated hydrogen sulfide on human airways and associated immune cells: Implications for respiratory diseases. *Front Public Heal*. 2019;7(JUN).
 91. Li L, Liu Y, Wang Q, Wang Z, Cui L, Xu Y, et al. Levels of nasal exhaled hydrogen sulfide in the general population and allergic rhinitis patients. *J Clin Lab Anal*. 2021;35(3):1–7.
 92. Perry MM, Tildy B, Papi A, Casolari P, Caramori G, Rempel KL, et al. The anti-proliferative and anti-inflammatory response of COPD airway smooth muscle cells to hydrogen sulfide. *Respir Res*. 2018;19(1):1–10.
 93. Khattak S, Zhang Q-Q, Sarfraz M, Muhammad P, Ngowi EE, Khan NH, et al. The Role of Hydrogen Sulfide in Respiratory Diseases. *Biomolecules*. 2021;11(5):682.
 94. Citi V, Martelli A, Brancaleone V, Brogi S, Gojon G, Montanaro R, et al. Anti-inflammatory and antiviral roles of hydrogen sulfide: Rationale for considering H₂S donors in COVID-19 therapy. *Br J Pharmacol*. 2020;177(21):4931–41.
 95. Hamidzadeh N, Simaetabar S, Handjani F, Ranjbar S, Moghadam MG PM. Composition of minerals and trace elements at Mamasani thermal source: A possible eventive treatment for some skin diseases. *J Edu Heal Promot*. 2017;6(110):1–6.
 96. Cacciapuoti S, Luciano M, Megna M, Annunziata M, Napolitano M, Patruno C, et al. The Role of Thermal Water in Chronic Skin Diseases Management: A Review of the Literature. *J Clin Med*. 2020;9(9):3047.
 97. José J, Galve G, Saz Peiró P, Lucas MO, Torres AH, Gil ES, et al. Quality of life and assessment after local application of sulphurous water in the home environment in patients with psoriasis vulgaris: A randomised placebo-controlled pilot study. *Eur J Integr Med* [Internet]. 2012;4:213–8. Available from: www.sciencedirect.com
 98. Carubbi C, Gobbi G, Bucci G, Gesi M, Vitale M, Mirandola P. Skin, inflammation and sulfurous waters: What is known, what is believed. *Eur J Inflamm*. 2013;11(3):591–9.
 99. Merighi S, Gessi S, Varani K, Fazzi D, Borea PA. Hydrogen sulfide modulates the release of nitric oxide and VEGF in human keratinocytes. *Pharmacol Res* [Internet]. 2012;66:428–36. Available from: <http://dx.doi.org/10.1016/j.phrs.2012.07.002>
 100. Xiao Q, Xiong L, Tang J, Li L, Li L. Hydrogen Sulfide in Skin Diseases: A Novel Mediator and Therapeutic Target. *Oxid Med Cell Longev*. 2021;2021.
 101. Mirandola P, Gobbi G, Micheloni C, Vaccarezza M, Di Marcantonio D, Ruscitti F, et al. Hydrogen sulfide inhibits IL-8 expression in human keratinocytes via MAP kinase signaling. *Lab Investig*. 2011;91(8):1188–94.
 102. Gross-Amat O, Guillen M, Gimeno JP, Salzet M, Lebonvallet N, Misery L, et al. Molecular mapping of hydrogen sulfide targets in normal human keratinocytes. *Int J Mol Sci*. 2020;21(13):1–14.
 103. Carson RJ, Konje JC. Role of hydrogen sulfide in the female reproductive tract. *Expert Rev Obstet Gynecol*. 2010;5(2):203–13.
 104. Lei S, Cao Y, Sun J, Li M, Zhao D. H₂S promotes proliferation of endometrial stromal cells via activating the NF- κ B pathway in endometriosis. *Am J Transl Res*. 2018;10(12):4247–57.
 105. Subadi I, Nugraha B, Laswati H, Josomuljono H. Pain relief with wet cupping therapy in rats is mediated by heat shock protein 70 and β -endorphin. *Iran J Med Sci*. 2017;42(4):384–91.
 106. Costa SKPF, Muscara MN, Allain T, Dallazen J, Gonzaga L, Buret AG, et al. Enhanced analgesic effects and gastrointestinal safety of a novel, hydrogen sulfide-releasing anti-inflammatory drug (atb-352): A role for endogenous cannabinoids. *Antioxidants Redox Signal*. 2020;33(14):1003–9.
 107. Wallace JL, Nagy P, Feener TD, Allain T, Ditrói T, Vaughan DJ, et al. A proof-of-concept, Phase 2 clinical trial of the gastrointestinal safety of a hydrogen sulfide-releasing anti-inflammatory drug. *Br J Pharmacol*. 2020;177(4):769–77.
 108. Kloesch B, Liszt M, Krehan D, Broell J, Kiener H, Steiner G. High concentrations of hydrogen sulphide elevate the expression of a series of pro-inflammatory genes in fibroblast-like synoviocytes derived from rheumatoid and osteoarthritis patients. *Immunol Lett*. 2011;141:197–203.